

BEDROCK GEOLOGIC MAP OF THE SALEM (IOWA) 7.5' QUADRANGLE

LEGEND

CENOZOIC

QUATERNARY SYSTEM

Qu - Undifferentiated Unconsolidated Sediments - Consists of heavy silty clay with developed loess, glacial till, and colluvium of variable thickness, and alluvial fans, silt, sand, and gravel. The total thickness of the Quaternary deposits varies between 0 ft (0 m) - 60 ft (18 m), but can be as much as 20 m (130 ft) thick in the eastern part of the mapping area. This unit is shown only on the cross-section, not on the map.

PALEOZOIC

CARBONIFEROUS SYSTEM

PENNSYLVANIAN SUBSYSTEM

Pcl - **Shale and Sandstone** (Lower Cherokee Group) - Pennsylvanian strata occur as scattered outcrops reaching a thickness of up to 13 m (43 ft) within the mapping area. This unit consists of shale, sandstone, light to medium gray, silty, sandy, and fine to medium grained sandstone, mostly argillaceous. Some shales are carbonaceous to pyrobituminous, with part coal. No outcrop of this unit was identified in the mapping area.

MISSISSIPPIAN SUBSYSTEM

Mpsl - **Limestone, Sandstone, and Dolomite** (Pella or "St. Louis" Formation) Middle Upper Mississippian, Meramec lower Ozarkian. The map area ranges from 91° 13' W to 91° 30' W and includes a maximum thickness of 20 m (66 ft) in the mapping area. It is dominated by limestone, sandstone, dolomite limestone, and dolomite with minor shale and silt. Limestone of the Pella Formation is typically white to light gray with scattered abundant fossils, generally brachiopods, trilobites, and corals. The "St. Louis" Formation is dominated by limestone, sandy limestone, sandstone, and dolomite, variably cherty. The limestone facies of this unit can be facies with brachiopods, trilobites, and several species of coral while the dolomite facies typically exhibit fossiliferous. Some fossils are encrusted. Sandstones of the "St. Louis" Formation are typically very fine to medium grained sandstone that are poorly to moderately cemented with calcareous cement. The lower portion of the "St. Louis" Formation commonly gray to dark brown dolomite, locally brecciated and sandy, with rare fossils. This mapping unit delineates the bedrock surface in the mapping area and is overlain by Quaternary sediments or Pennsylvanian rocks. No outcrop was identified in the mapping area although scattered exposures of the lower portion of this mapping unit was observed in a small quarry in the extreme western corner of the mapping area.

Mws - **Shale, Dolomite, and Limestone** (Warsaw Formation) Upper Ozarkian. The Warsaw Formation varies in thickness reaching a maximum thickness of approximately 17 m (56 ft). This unit is generally bedded into two major lithologic groups, a lower argillaceous dolomite sequence and an upper shale dominated sequence. The upper shale is typically light to medium gray, silty, and variably dolomitic with minor chert, sand, and sparse quartz grains. The lower dolomite, sometimes referred to as the "granite beds", is argillaceous to shaly, with scattered abundant quartz grains. Minor limestone outcrops occur locally in this unit, usually beds with crinoidal packstone grainstone facies. Brachiopods, trilobites, and bryozoans are found throughout the mapping area, although these fossils are more common in the carboniferous lithologies. This unit exhibits wide variability, having only the upper shale or lower dolomite facies, suggesting unusual depositional environments. Outcrops of this unit were not observed in the mapping area.

Mkeo - **Limestone, Sandstone, Chert, and Shale** (Keokuk Formation) Upper Ozarkian. The Keokuk Formation typically ranges from 2.2 m (7.2 ft) to 10 m (33 ft) in thickness in the mapping area. This unit is subdivided into two major lithologic groups, a lower argillaceous dolomite sequence and an upper shale dominated sequence. The upper shale is typically light to medium gray, silty, and variably dolomitic with minor chert, sand, and sparse quartz grains. The lower dolomite, sometimes referred to as the "granite beds", is argillaceous to shaly, with scattered abundant quartz grains. Minor limestone outcrops occur locally in this unit, usually beds with crinoidal packstone grainstone facies. Brachiopods, trilobites, and bryozoans are found throughout the mapping area, although these fossils are more common in the carboniferous lithologies. This unit exhibits wide variability, having only the upper shale or lower dolomite facies, suggesting unusual depositional environments. Outcrops of this unit were not observed in the mapping area.

Mb - **Limestone, Dolomite, and Chert** (Burlington Formation) Lower Ozarkian. The Burlington Formation can be up to 20 m (66 ft) thick in the mapping area. This unit is subdivided into three members (in ascending order) the Debar Creek, Hagler Creek, and Cedar Fork, characterized by distinct lithologic groupings. The Debar Creek Member is dominated by white to light gray limestone displaying packstone grainstone facies and nodular fossiliferous chert. The Hagler Creek Member is characterized by dolomite with an associated sand and siltstone facies. The Cedar Fork Member is a thin bedded, argillaceous to shaly dolomite. The Burlington Formation is a high to medium grained, dolomite limestone that grades to an argillaceous to shaly dolomite. The Burlington Formation is a high to medium grained, dolomite limestone that grades to an argillaceous to shaly dolomite. The Burlington Formation is a high to medium grained, dolomite limestone that grades to an argillaceous to shaly dolomite.

Mk - **Dolomite, Limestone, and Silstone** (Kinderhook Formation) Lower Mississippian. The Kinderhook sequence ranges in thickness from 15 m (49 ft) to 20 m (66 ft) with a maximum thickness of 27 m (89 ft) in the mapping area. This unit comprises three formations in ascending order: the McCroney, Prospect Hill, and Warsaw, characterized by distinct lithologic groupings. The McCroney Formation is composed of alternating beds of argillaceous limestone, sub-lithologic limestone and dark brown, argillaceous dolomite generating a unique "mudstone" appearance on outcrop. A hard, often locally present. The Prospect Hill Formation is a high to medium grained, dolomite limestone that grades to an argillaceous to shaly dolomite. The Warsaw Formation is a high to medium grained, dolomite limestone that grades to an argillaceous to shaly dolomite.


DEVONIAN SYSTEM

Der - **Siltstone and Shale** (English River Formation) Upper Devonian. The English River Formation can be up to 6 m (20 ft) thick in the mapping area. This unit is dominated by gray to green silty shale with argillaceous dolomite. Dolomite and brachiopods or corals are typically in the upper beds, with siltstone to abundant fossiliferous shale. This unit is only present in the extreme western corner of the mapping area.

Dss - **Shale** (Saverton Shale Formation) Upper Devonian. The Saverton Shale Formation can be up to 10 m (33 ft) thick in the mapping area. This unit is dominated by gray to green shale, commonly barren with sparse to absent macrofossils. This unit only appears in the cross-section, not on the map.


OTHER FEATURES

- New drill holes for this map project
- New geophysical data point
- Geological Survey data points
- Geological Survey data points - records available at www.iowageologicalsurvey.org
- Overlays
- Incorporated towns or cities
- Roads
- Well used for geologic cross-section
- Bedrock thickness - shades of gray show the bedrock surface as it would be illuminated by an artificial light source from the NW direction.




Adjacent 7.5' Quadrangles

LOCKRIDGE EAST, IOWA	MOUNT PLEASANT, IOWA	NEW LODON, IOWA
HILLSBORO, IOWA	SALEM, IOWA	LOWHILL, IOWA
FARMINGTON, IOWA	CONNELLSON, IOWA	WEST POINT, IOWA



STRATIGRAPHIC COLUMN



BEDROCK GEOLOGIC MAP OF THE SALEM 7.5' QUADRANGLE, HENRY AND LEE COUNTIES, IOWA

IOWA GEOLOGICAL SURVEY
OPEN FILE MAP OFM-19-3
JUNE 2019

Ryan Clark, Stephanie Tasser-Satine, and Phil Kerr
Iowa Geological Survey, IHR-Hydroscience & Engineering, University of Iowa, Iowa City, Iowa

IOWA GEOLOGICAL SURVEY
Iowa Geological Survey, Keith Schilling, State Geologist

Supported in part by the U.S. Geological Survey, National Cooperative Geologic Mapping Program (NCEM/NAMP). This map is partially funded by a National Science Foundation grant (NSF-EAR-160629) through the Iowa State University STEM Education Grant (IMPACT-160629).

ACKNOWLEDGMENTS

Special thanks to the Smith of Crawford Commission, Co. (Iowa) Materials for allowing us to access aquifers and around the mapping area during field activities. University of Iowa (UI) Department of Earth and Environmental Sciences (DEES) and the Center for Earth and Environmental Science (CEES) provided well-coring support. Tom Schilling, geologic site manager for UI student Megan Koch and Athena Kambou, by providing downcore logs of water well drilling samples. Megan Koch and Athena Kambou also helped with well locations and data collection. Tom Schilling, State Geologist, Iowa Geological Survey (IGS) helped with geophysical data acquisition. Thanks also to Rick Langley of the IGSS for mapping the lower St. Louis Formation (Pella or "St. Louis" Formation) in Henry and Lee Counties. Professor of Geology, Iowa State University, Dr. Arthur H. Rowley, provided the lithologic column and lithologic symbols for the stratigraphic column components appearing on this mapping project. Casey Kerr and Chris Kahl of the Iowa Department of Natural Resources (IDNR) provided the well logs for the Skunk River and Skunk River area. The Iowa Department of Natural Resources (IDNR) provided the well logs for the Skunk River and Skunk River area. The Iowa Department of Natural Resources (IDNR) provided the well logs for the Skunk River and Skunk River area. The Iowa Department of Natural Resources (IDNR) provided the well logs for the Skunk River and Skunk River area.

INTRODUCTION TO THE BEDROCK GEOLOGY OF THE SALEM 7.5' QUADRANGLE, HENRY AND LEE COUNTIES, IOWA

The Salem Quadrangle lies within the northern Iowa Drift Plain landform region (Prior, 1991). The map area is dominated by loess-mantled till plains in the uplands, and fine to coarse grained alluvial deposits within the Skunk River and its tributaries. This area hosts glacial deposits of Pre-Illinoian age (ranging from 0.5 to 2.6 million years ago). The thickness of Quaternary materials overlying the bedrock surface varies widely across the quadrangle ranging from 0 to 18 m (0 - 60 ft), reaching a maximum thickness of 50 m (165 ft) within the bedrock valley in the northeastern part of the mapping area. Shallow bedrock information from the soil surveys of Henry and Lee counties (Scholten, 1965; and Lockridge, 1979) was used for identifying potential bedrock outcrop locations during field mapping activities. Bedrock outcrops were not observed during field activities, however according to the soil survey maps, outcrops exist along tributaries of the Skunk River, and in areas of the southeastern part of the quadrangle. There is one operational quarry in the extreme southeastern corner of the quadrangle that was observed during field mapping activities, as well as one abandoned quarry that was not accessible. Subsurface information was mostly derived from the analysis of water well cutting samples deposited at the Iowa Geological Survey (IGS). Lithologic and stratigraphic information from these samples are stored in the online GeoSam database of the IGS. Geologic information from one quarry and more than 250 private and public wells within the Salem Quadrangle and the surrounding area were used for bedrock geologic mapping purposes.

The Mississippian System (now Subsystem) was historically proposed for the succession of strata exposed in the Mississippi River Valley between Burlington, Iowa and southern Illinois. Therefore, the bedrock exposures in southeastern Iowa take on a special significance as they comprise part of the historic "body stratotype" on which the concept of the Mississippian System was defined and based (Witzke et al., 2002). The Mississippian had been primarily a North American chronostratigraphic label roughly synonymous with the Lower Carboniferous of the Old World. After approval by the Subcommittee on Carboniferous Stratigraphy in 1999 and ratification by the International Union of Geological Sciences and the International Commission on Stratigraphy in 2000, the Carboniferous System was officially subdivided into the lower and upper subsystems, the Mississippian and Pennsylvanian, respectively. As such, the Mississippian now has meaning and application as a major subdivision of geologic time not only in North America, but as a globally defined subsystem. The bedrock strata seen in the Salem Quadrangle and surrounding area provide a significant historic reference for the Mississippian as a whole.

The contention that is the Mississippian in Iowa has been the subject of curiosity for many previous workers. Owen (1852) and Hall (1857) were the first to recognize that the abundant bedrock exposures in southeastern Iowa likely correlated with those observed farther down the Mississippi River Valley. Decades later, Van Tuyl (1923) took on the ambitious task of correlating all of the Mississippian units across Iowa. Many of their lithologic interpretations were valuable; however, the correlations were, and continue to be, subject to revision as later workers attempted to piece the Mississippian into the global stratigraphic framework. Harris and Parker (1964) provided inspirational insights into the structural context of southeastern Iowa by identifying a series of northeast-southwest trending anticlines that were later found to be superimposed on the larger northeast-southwest trending structural feature known as the Mississippi Arch (Witzke et al., 1990). Many questions remain regarding the stratigraphic correlations within the Mississippian such as whether the "St. Louis" Formation in Iowa truly belongs in the St. Louis Formation or whether some of the upper members be reassigned to the St. Genevieve Formation; whether the Prospect Hill Formation is an offshoot of the Hannibal Formation of Missouri and Illinois; and whether the McCroney Formation is correlative to the McCroney in Illinois or if it should become a new stratigraphic interval (as proposed by Witzke et al., 2002). In an effort to address the question regarding the "St. Louis" Formation, detrital zircon analyses from sandstone samples collected near the mapping area were processed with the help of Emily Fritzel (Assistant Professor of Geology at the University of Iowa (UI)). The geochronologic data provided by the detrital zircon analysis were not able to differentiate the sandstone units within the "St. Louis" Formation, however, further study of the geochemistry and lithology of these sandstones may provide the evidence needed to identify whether these units belong in the St. Louis proper or in the St. Genevieve. Clarification of the issue regarding the Prospect Hill and McCroney formations is being carried out with the help of Brad Crane (Assistant Professor of Geology at the UI), Brittany Stofius (UI student), and James "Jed" Day (Professor of Geology at Illinois State University). Samples collected from locations southeast of the mapping area as well as other locations in eastern Illinois and northeastern Missouri for conodonts and carbon isotopes have provided valuable bio- and chemostratigraphic information. Preliminary results suggest that the Prospect Hill and McCroney formations in Iowa may correlate with the Hannibal Formation of Missouri and Illinois. Further study will commence with additional sampling of surface exposures as well as core samples. Rectifying the questions posed by Witzke et al., 2002, may now become attainable.

Although the Mississippian bedrock in southeastern Iowa is no longer a widely used aquifer due to low yields and locally poor water quality, many of the bedrock units are highly desirable sources of aggregate, thus necessitating the continued effort to gain a better understanding of the local and regional stratigraphic characteristics and relationships of the Mississippian Subsystem in southeastern Iowa. The mapping area consists of bedrock of the Mississippian Subsystem from late Kinderhookian to early Chesterian (about 255 - 330 million years ago) and Devonian strata of Famennian age (about 370 - 360 million years ago) (Ogg et al., 2008). Famennian strata are represented by brown, organic rich shales of the Grassy Creek Formation overlain by gray-green silty shales of the Saverton Shale Formation and capped by the English River Formation siltstone. The thick shale packages represent major transgressive-regressive cycles of deposition in a stratified seaway (Witzke, 1987). Kinderhookian strata represent a sequence of interbedded carbonates and siltstones that unconformably underlie the Burlington Formation (early Ozarkian) within the mapping area. The Burlington, Keokuk, and Warsaw formations (collectively the Augusta Group of Witzke et al., 2010) represent a relatively conformable package of marine rocks deposited during the Ozarkian transgressive-regressive (T-R) cycle. Interpreted as part of the central middle shelf of the Ozarkian sea that transgressed toward the northwest and the Transcontinental Arch, the Burlington Formation rocks were deposited on a vast subtidal epicontinental shelf that stretched from Illinois and Iowa into central Kansas and Oklahoma (Lane, 1978; Witzke et al., 1990). The Keokuk and Warsaw formations represent the regressive phase of the Ozarkian T-R cycle punctuated by a stark unconformity below the overlying Pella or "St. Louis" formations, regionally displaying up to 40 m (130 ft) of erosional relief (Witzke et al., 2002). For a more detailed description of the lithologic units and further discussion of mapping methodologies, please refer to the accompanying Summary Report.

REFERENCES

Hall, J., 1857. Observations upon the Carboniferous limestones of the Mississippi Valley. American Journal of Science, v. 23, p. 187-203.

Harris, G. S., and Parker, M. C., 1964. Stratigraphy of the Ozage Series in Southeast Iowa, Iowa Geological Survey Report of Investigation No. 1, 52 p., 17 Figs., 7 plates.

Lane, H. R., 1978. The Burlington Shelf (Mississippian, north-central United States). Geologica et Palaeontologica v. 12, p. 165-176.

Lockridge, L. D., 1979. Soil Survey of Lee County, Iowa. U.S. Department of Agriculture, Soil Conservation Service, 188 p., 85 map sheets.

Ogg, J. G., Ogg, G., and Gradstein, F. M., 2008. The Concise Geologic Time Scale, Cambridge University Press, 177 p.

Owen, D. D., 1852. Report of a Geological Survey of Wisconsin, Iowa, and Minnesota, and incidentally of a portion of Nebraska territory. Lippincott, Gambo and Co., 638 p.

Prior, J. C., 1991. Landforms of Iowa. Univ. of Iowa Press, Iowa City, 154 p.

Scholten, J. E., 1965. Soil Survey of Henry County, Iowa. U.S. Department of Agriculture, Soil Conservation Service, 259 p., 63 map sheets.

Van Tuyl, F. M., 1923. The stratigraphy of the Mississippian formations of Iowa. Iowa Geological Survey, Annual Report, v. 30, p. 33-59.

Witzke, B. J., 1987. Models for circulation patterns in epicontinental seas applied to Paleozoic facies of North American craton. Paleogeography, v. 2, p. 229-248.

Witzke, B. J., Anderson, R. R., and Pope, J. P., 2010. Bedrock Geologic Map of Iowa, 1:500,000. Iowa Geological and Water Survey Open File Map OFM-10-1.

Witzke, B. J., Banker, B. J., Anderson, R. R., Ariz, J. A., and Tasser-Satine, S. A., 2002. Pleistocene, Mississippian, & Devonian Stratigraphy of the Burlington, Iowa, Iowa Geological Survey Guidebook No. 23, 137 p.

Witzke, B. J., McKay, R. M., Banker, B. J., and Woodson, F. J., 1990. Stratigraphy and Paleogeography of Mississippian Strata in Keokuk and Washington Counties, Southeast Iowa. Iowa Geological Survey Guidebook No. 10, 105 p.

Base map from USGS Salem 7.5' Digital Raster Graphic (DSG) file, IA_Salem_USGS_30m.tif which was scanned and modified from the Salem 7.5' Topographic Quadrangle map, published by the US Geological Survey in 2015. Land elevation contours (1' interval). Bedrock topography raster created internally for this map project.

Iowa Geological Survey digital cartographic file: Salem_BedrockGeology.mxd, version 6/30/19 (AUG19 10:5).

Map projection and coordinate system based on Universal Transverse Mercator (UTM) Zone 19A, datum NAD83.

The map and cross-section are based on interpretations of the best available information at the time of mapping. Map interpretations are not a substitute for detailed site specific studies.

Research supported by the U.S. Geological Survey, National Cooperative Geologic Mapping Program, under USGS award number 01AC020198. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

GEOLOGIC CROSS-SECTION A-A'

