

## INTRODUCTION

Two 1:24,000 quadrangle scale surficial maps were produced as part of a multi-phase project in developing areas of Scott County. The Dixon quadrangle covers an area from 41° 37' 30" to 41° 45' N latitude and 90° 45' to 90° 52' 30" W longitude. The Eldridge quadrangle covers an area from 41° 37' 30" to 41° 45' N latitude and 90° 30' to 90° 37' 30" W longitude. The mapping area is dominated by the Southern Iowa Drift Plain landform region, but includes the boundary of the Iowan Erosion Surface landform region in the northern part of the Eldridge quadrangle. The area has been subjected to multiple periods of Quaternary glaciations and subaerial erosion. The mapping region includes areas glaciated during the Illinois and Pre-Illinois episodes. Periodic erosion during the last 500,000 years has led to the development of an integrated drainage network and the destruction of pre-existing glacial landforms associated with Illinoian and Pre-Illinoian glaciations. Generally speaking, the map area consists of Wisconsin age loesses mantling Illinoian and Pre-Illinoian glacial sediments of variable thickness. These deposits are regionally extensive. Previous surficial geologic mapping of the map area consists of the Des Moines 4° x 6° Quadrangle at a scale of 1:1,000,000 (Hallberg et al., 1991). The prior STATEMAP project, Surficial Geologic Materials of the McCausland Quadrangle (Quade et al., 2004) is adjacent to the current map area.

## BRIEF GEOLOGIC HISTORY

Early researchers believed there were only two episodes of Pre-Illinoian glaciation in Iowa: Kansan and Nebraskan. Later regional studies determined that at least seven episodes of Pre-Illinoian glaciation occurred in this region from approximately 2.2 million to 500,000 years ago (Boellstorff, 1978a; Boellstorff 1978b; Hallberg, 1980a; Hallberg, 1986). Hallberg (1980a, 1980b, 1986) undertook a regional scale project that involved detailed outcrop and subsurface investigations including extensive laboratory work and synthesis of previous studies. This study led to the abandonment of the classic glacial and interglacial terminology: Kansan, Aftonian and Nebraskan. Hallberg's study marked a shift from use of time-stratigraphic terms to lithostratigraphic classification. The result of Hallberg's study was the development of a lithostratigraphic framework for Pre-Illinoian till. In east-central Iowa, Hallberg formally classified the units into two formations on the basis of differences in clay mineralogy: the Alburnett Formation (several undifferentiated members) and the younger Wolf Creek Formation (including the Winthrop, Aurora and Hickory Hills members). Both formations are composed predominantly of till deposits, but other materials are present. Paleosols are formed in the upper part of these till units.

Following the Pre-Illinoian glaciations, most areas of eastern Iowa underwent extensive landscape development and erosion until the Wisconsin Episode loess began to be deposited. However, a limited area of southeastern Iowa was glaciated during the Illinois Episode, around 300,000-130,000 years ago. The majority of the mapping area in Scott County was glaciated during this time; however, the area north of Mud Creek was not glaciated. Leverett (1898, 1899) was the first to study the Illinoian glacial deposits in southeast Iowa. The Illinoian till was deposited by the advancing Lake Michigan Lobe which moved across western Illinois into Iowa from the northeast (Leverett, 1899; Wickham, 1980). The Lake Michigan Lobe incorporated Paleozoic bedrock materials from the Lake Michigan Basin which are distinguished by both the clay mineralogy of the matrix as well as the pebbles and clasts in the Illinoian deposits (Lineback, 1980; Wickham, 1980). Till fabrics and glacial landform orientations also provide evidence of a northeastern source area for these tills (Lineback, 1979). The differing provenances between the Illinoian and Pre-Illinoian tills (which moved into Iowa from north-northeast) create identifiable and distinguishing physical and mineralogical characteristics. Subsequent researchers have made few changes to the original concept of the Illinoian stage as defined by Leverett. The most significant change was the subdivision into three substages (Willman et al., 1963; Frye et al., 1964; Willman and Frye, 1970). Work during the 1950's and 1960's focused on characterizing the mineralogy and petrography. The Pleistocene

deposits and stratigraphy of Illinois were more formally classified by Willman and Frye in 1970, and the oldest Illinoian age till was named the Kellerville Till Member of the Glasford Formation. Hallberg (1980b) defined the formal stratigraphic nomenclature in Iowa. The only Glasford Formation till present in Iowa is the Kellerville Till Member which was described from roadcut exposures in western Illinois (Willman and Frye, 1970). The type section is located in southeastern Adams County, Illinois, approximately 100 km (62 miles) south of Burlington, Iowa. The Kellerville Till Member replaces the terms Mendon Till (Frye et al., 1964; Frye et al., 1969) and Payson Till (Leighton and Willman, 1950; Wanless, 1957).

In eastern Iowa, the highly eroded and dissected Illinoian and Pre-Illinoian upland and older terraces are mantled by Wisconsin loesses of variable thickness (Ruhe, 1969; Prior, 1976). These sediments are the youngest regionally extensive Quaternary deposits and were deposited between 30,000 and 12,000 years ago. Loess is thickest in the region near the Iowan Erosion Surface (IES) boundary and near local sources (Wapsipinicon River valley). Two loess units were deposited across eastern Iowa, the older Pisgah Formation and the younger Peoria Loess. The Pisgah is thin and includes loess and related slope sediments that have been altered by colluvial hillslope processes, pedogenic and periglacial processes. The unit is characterized by the presence of a weakly developed soil recognized as the Farmdale Geosol. It is not uncommon to see the Farmdale developed throughout the Pisgah and incorporated into the underlying older Sangamon Geosol. Most likely the Pisgah loess was deposited on the eastern Iowa landscape from 30,000 to 24,000 years ago (Bettis, 1989). The Pisgah Formation is typically buried by Peoria Formation loess. The Peoria Formation loess accumulated on stable landsurfaces in eastern Iowa from 25,000 to 21,000 years ago and was followed by a period of intense cold during the Wisconsin full glacial episode from 21,000 to 16,500 years ago (Bettis, 1989). Outside the IES, Peoria Loess continued to accumulate until 13,000 B.P.; and in some parts of the IES a thin increment of loess accumulated as the climate ameliorated approximately 14,000 to 12,000 years ago. This period of intense cold and ensuing upland erosion led to the development of the distinctive landform recognized as the IES (Prior, 1976). During this period surface soils were removed from the IES and the Pre-Illinoian till surface was significantly eroded; resulting in the development of a lag deposit referred to as a "stone line". The depositional history of the IES was under great debate for an extended period of time. Early researchers believed the IES was a separate glaciation occurring sometime between the Illinois and the Wisconsin episodes. Later work disproved this idea and determined that erosional processes controlled the landscape development (Ruhe et al., 1968). Hallberg et al. (1978) revisited the "Iowan Erosion Surface" to further research studies into the mechanisms behind the formation of the erosion surface and to reiterate Ruhe's classic work and to illustrate the need for continued research in the area.

Esling (1984) undertook a regional study to document extensive post Illinoian-age alluvial deposits that had accumulated in major valleys in eastern Iowa. Three major terrace assemblages with differing stratigraphy and age were identified: Early Phase High Terrace (EPHT), Late Phase High Terrace (LPHT) and Low Terrace (LT). EPHT deposits are characterized by the presence of Peoria and Pisgah Formation sediments overlying a Sangamon Geosol in the underlying alluvium. Esling theorized that these terraces are older than 40,000 years B.P. but younger than the Illinoian sediments in eastern Iowa (Bettis, 1989). LPHT deposits are characterized by the presence of Peoria Formation loess grading down into underlying alluvium with no paleosol. These terraces developed prior to 25,000 years ago and were buried by loess before 12,500 years ago. The LPHT terraces are typically inset into EPHT deposits. The LT is the youngest terrace and is not buried by Peoria loess. In the map area, eolian dunes (Peoria Formation-sand facies) are present on the terrace surface and indicate this terrace surface was deposited during Late Wisconsin through early Holocene time (12,500 to 10,000 years ago).

The pre-Holocene high terrace identified by Esling (1984) in the Iowa and Cedar River valleys has a similar stratigraphic sequence as the deposits along Mud Creek in Scott County. Autin and Bettis (1991) undertook an extensive study of the development of Mud Creek in eastern Iowa. One of the project goals

was to define a stratigraphic and geomorphic framework for the region. Four mapping units were identified: an upland erosion surface, a high terrace, an intermediate terrace level, and the Mud Creek floodplain deposits. The upland erosion surface consists of the Yarmouth-Sangamon, Late Sangamon, and Iowan erosion surface materials overlain by Peoria Formation loess. Pisgah Formation materials may be present below the Peoria Formation and appears to be an upland pedisegment facies related to the Late Wisconsin alluvium. In the northwestern part of the Dixon quadrangle, a dune field of Late Wisconsin sand is buried by the Peoria Loess in broad upland positions. The high terrace unit consists of Peoria Formation Loess and Pisgah Formation overlying Pre-Wisconsinan alluvium, with a paleosol developed at the top of the alluvium. This unit is analogous to the Early Phase High Terrace described by Esling (1984). The distribution suggests the relationship with the aggradation of a river in the Cleona Channel (Autin and Bettis, 1991). The intermediate terrace is associated with Late Wisconsin constructional surfaces. This surface consists of Peoria Loess overlying alluvium. The alluvium may be underlain by Pisgah Formation, Pre-Illinoian till or bedrock. This unit occurs on the lower upland surface along the Mud Creek floodplain as an erosional remnant. Mud Creek floodplain deposits include the Camp Creek, Roberts Creek, Gunder, and Pre-Gunder members of the Holocene DeForest Formation.

## **DESCRIPTION OF LANDFORM SEDIMENT ASSEMBLAGE MAP UNITS**

Recent studies and mapping indicate that the map area encompasses a complex suite of depositional landforms and sediment sequences related to glaciations, alluviation, subaerial erosion, and wind-blown transport. To map diverse landscapes we have selected the most comprehensive mapping strategy—a landform sediment assemblage (LSA) approach. Various landforms are the result of specific processes at work in the geologic system. Landforms typically have similar relief, stratigraphic and sedimentologic characteristics. Recognition of the genetic relationship among landforms and their underlying sediment sequences allows one to generalize and map complex glacial terrains over areas of large extent (Sugden and John, 1976; Eyles and Menzies, 1983). Bettis and others (1999) found LSA mapping concepts were extremely useful in overcoming the difficulties of mapping in large valleys and noted LSA's provided a unique opportunity to associate landforms with their underlying sediment packages.

Thirteen landform sediment assemblage units were identified in the map area utilizing orthophotos, topographic expression, digitized soil maps and existing and new subsurface boring information. Stratigraphic information from Mud Creek (Autin and Bettis, 1991) proved very useful in mapping the extent of Holocene and Wisconsin age terraces in the map area. The thirteen LSA units are: Hudson Episode: Alluvium, Thick Alluvium; Wisconsin Episode: Late Phase High Terrace (LPHT), Early Phase High Terrace (EPHT), Loess over Glasford Formation, Thick Loess over Glasford Formation, Loess and Intercalated Eolian Sand over Glasford Formation, Thick Loess and Intercalated Eolian Sand over Glasford Formation, Loess over Pre-Illinoian till, Loess and Intercalated Eolian Sand over Pre-Illinoian till, Loamy and Sandy Sediment Shallow to Glacial Till; Illinois Episode: Glacial Till; and Pre-Illinois Episode: Glacial Till. The following is a description of each landform sediment assemblage listed in order of episode.

### **HUDSON EPISODE**

#### **Landform Sediment Assemblages**

**Qal - Alluvium (DeForest Formation-Undifferentiated)** Variable thickness (<1 to 5 meters) of very dark gray to brown, noncalcareous to calcareous, stratified silty clay loam, clay loam, loam to sandy loam alluvium and colluvium in stream valleys, on hill slopes and in closed depressions. May overlie Glasford Formation glacial till or Noah Creek Formation. Associated with low-relief modern floodplain, closed

depressions, modern drainageways or toeslope positions on the landscape. Seasonal high water table and potential for frequent flooding.

**Qa2 - Thick Alluvium (DeForest Formation-Undifferentiated)** Two to five meters of massive to moderately well stratified loam, silt loam, clay loam, or loamy sand overlying five to thirty meters of poorly to moderately well sorted, massive to well stratified, coarse to fine feldspathic pebbly sand and gravel of the Noah Creek Formation. This unit locally serves as an aquifer. Seasonally high water tables occur in this map unit.

## WISCONSIN EPISODE

### Landform Sediment Assemblages

**Qptlp - Late Phase High Terrace (LPHT) (Peoria Formation-silt and/or sand facies)** Two to seven meters of yellowish brown to gray, massive, jointed, calcareous or noncalcareous, silt loam and intercalated fine to medium, well sorted, sand. Grades downward to poorly to moderately well sorted, moderately to well stratified, coarse to fine feldspathic quartz sand, loam, or silt loam alluvium.

**Qptep - Early Phase High Terrace (EPHT) (Peoria Formation-silt and/or sand facies)** Two to seven meters of yellowish brown to gray, massive, jointed, calcareous or noncalcareous, silt loam and intercalated fine to medium, well sorted, sand. The Peoria deposits overlie a Farmdale Geosol developed in the Pisgah Formation which in turn overlies a well-expressed Sangamon Geosol developed in poorly to moderately well sorted, moderately to well stratified, coarse to fine sand, loam, or silt loam alluvium.

**Qps\_gla - Loess (Peoria Formation-silt facies)** Generally 2 to 5 m of yellowish to grayish brown, massive, jointed calcareous or noncalcareous silt loam to silty clay loam. Overlies a grayish brown to olive gray silty clay loam to silty clay (Pisgah Formation—eroded Farmdale Geosol) which is less than 1.5 m thick. The Farmdale Geosol appears to be disturbed by periglacial action and is welded to an older Sangamon Geosol developed in loamy glacial till of the Glasford Formation. This mapping unit encompasses upland divides, ridgetops and convex sideslopes. Well to somewhat poorly drained landscape.

**Qpsb\_gla - Thick Loess (Peoria Formation—silt facies)** Generally 5 to 15 m of yellowish to grayish brown, massive, jointed calcareous or noncalcareous silt loam to silty clay loam. Overlies massive, fractured, clay loam glacial till of the Glasford formation with or without intervening clayey Farmdale Geosol. This mapping unit encompasses upland divides, ridgetops and convex sideslopes. Well to somewhat poorly drained landscape.

**Qps1\_gla - Loess and Intercalated Eolian Sand (Peoria Formation-silt facies)** Two to ten meters of yellowish brown to gray, massive, fractured, noncalcareous grading downward to calcareous silt loam and intercalated fine to medium, well sorted, sand. Sand is most abundant in lower part of the eolian package. Overlies massive, fractured, loamy glacial till of the Illinoian Glasford Formation till with or without intervening clayey Farmdale /Sangamon Geosol.

**Qps1b\_gla -Thick Loess and Intercalated Eolian Sand (Peoria Formation-silt facies)** – Five to fifteen meters of yellowish brown to gray, massive, noncalcareous grading downward to calcareous silt loam and intercalated fine to medium, well sorted, sand. Minimum thickness of five meters on uplands. Maximum thickness of two to seven meters of loess occurs on adjacent slopes. Overlies massive, fractured, loamy glacial till of the Illinoian Glasford Formation with or without intervening clayey Farmdale /Sangamon Geosol.

**Qps - Loess (Peoria Formation—silt facies)** Generally 2 to 8 m of yellowish to grayish brown, massive, jointed calcareous or noncalcareous silt loam to silty clay loam. Overlies a grayish brown to olive gray silty clay loam to silty clay (Pisgah Formation—eroded Farmdale Geosol) which is less than 1.5 m thick. The Farmdale Geosol appears to be disturbed by periglacial action and is welded to an older Sangamon Geosol developed in loamy glacial till of the Wolf Creek or Alburnett formations. This mapping unit encompasses upland divides, ridgetops and convex sideslopes. Well to somewhat poorly drained landscape.

**Qps1 - Loess and Intercalated Eolian Sand (Peoria Formation-silt facies)** Two to ten meters of yellowish brown to gray, massive, fractured, noncalcareous grading downward to calcareous silt loam and intercalated fine to medium, well sorted, sand. Sand is most abundant in lower part of the eolian package—eolian dunes 0.5 – 2.5 m thick may be present at the base of this unit. Overlies massive, fractured, loamy glacial till of the Wolf Creek or Alburnett formations with or without intervening clayey Farmdale/Sangamon Geosol.

**Qgla2 - Loamy and Sandy Sediment Shallow to Glacial Till (Unnamed erosion surface sediment )** One to three meters of yellowish brown to gray, massive to weakly stratified, well to poorly sorted loamy, sandy and silty erosion surface sediment. Map unit includes some areas mantled with less than two meters of Peoria Silt (loess). Overlies massive, fractured, firm glacial till of the Illinoian Glasford Formation. Seasonally high water table may occur in this map unit.

## ILLINOIS EPISODE

### Landform Sediment Assemblages

**Qgla - Till (Glasford Formation)** Generally 10 to 35 m of very dense, massive, fractured, loamy glacial till of the Illinoian Glasford Formation with or without a thin loess mantle (Peoria Formation—less than 2 meters) and intervening clayey Farmdale/ Sangamon Geosol. This mapping unit encompasses narrowly dissected interfluves and side slopes, and side valley slopes. Drainage is variable from well drained to poorly drained.

## PRE-ILLINOIS EPISODE

### Landform Sediment Assemblages

**Qwa3 – Till (Wolf Creek or Alburnett Formations)** Generally 10 to 35 m of very dense, massive, fractured, loamy glacial till of the Wolf Creek or Alburnett Formations with or without a thin loess mantle (Peoria Formation—less than 2 meters) and intervening clayey Farmdale/ Sangamon Geosol. This mapping unit encompasses narrowly dissected interfluves and side slopes, and side valley slopes. Drainage is variable from well drained to poorly drained.

## PALEOZOIC

**Su - Bedrock Exposures** (Silurian undifferentiated)

**Qf - Fill** Areas of major land filling. Fill associated with railroad grades, highway grades and land leveling. Variable in texture ranging from loamy to sandy to concrete rubble. Extent mapped as shown in county soil surveys.

**Qpq - Pits and Quarries** Sand and gravel pits and rock quarries. Extent mapped as shown in county soil surveys.

## **Water Features**

### **DESCRIPTION OF EAST-CENTRAL IOWA STRATIGRAPHY**

An important aspect of surficial geologic mapping on the Iowa Erosion Surface and the Southern Iowa Drift Plain is the development of map units that utilize previously established lithostratigraphic frameworks for the Hudson, Wisconsin, Illinoian and Pre-Illinoian deposits in Iowa. A stratigraphic framework allows us to better understand the surficial materials of east-central Iowa. Hudson, Wisconsin, Illinois and Pre-Illinois Episode deposits (Johnson et al., 1997) of east-central Iowa are included in six formations: DeForest Formation (Hudson), Peoria and Pisgah Formations (Wisconsin), Glasford Formation (Illinois) and Wolf Creek and Alburnett Formations (Pre-Illinois). The following section provides a description of formations and members of east-central Iowa deposits.

### **STRATIGRAPHIC FRAMEWORK FOR EAST-CENTRAL IOWA**

Surficial deposits of the map area are composed of six formations: DeForest, Peoria, Pisgah, Glasford, Wolf Creek, and Alburnett formations. Holocene age deposits associated with fine-grained alluvial and colluvial sediments include the DeForest Formation which is subdivided into the Camp Creek, Roberts Creek, Gunder, and Corrington members. The Peoria Formation includes wind-blown sediments: two facies are recognized, a silt facies (loess) and a sand facies (eolian sand). The Pisgah Formation originated as eolian silt that has been altered by a combination of colluvial hillslope processes, pedogenic and periglacial processes. The only Illinoian age deposits in Iowa consist of the Glasford Formation- Keller-ville Member. Pre-Illinoian glacial deposits in east-central Iowa consist of two formations: the younger Wolf Creek Formation and the Alburnett Formation. The Wolf Creek is divided into the Winthrop, Aurora and Hickory Hills members (oldest to youngest). The Alburnett Formation consists of several “undifferentiated” members.

### **DEFOREST FORMATION**

The DeForest Formation consists of fine-grained alluvium, colluvium, and pond sediment in stream valleys, on hillslopes and in closed and semi-closed depressions. The formation was originally defined by Daniels et al. (1963) for a repeatable sequence of alluvial fills in the Loess Hills of western Iowa. Subsequent study of drainage basins across Iowa revealed that a consistent alluvial stratigraphy was present, but its classification required expansion and revision of the formation (Bettis, 1990). The revised DeForest Formation includes the Gunder, Roberts Creek, Camp Creek, and Corrington members, all recognized on the DML (Bettis, 1990; Bettis et al., 1992). These members are not described here for the sake of brevity. These new members are the Flack Member, consisting of colluvium mantling hillslopes, the Woden Member, for sediment fills in semi-closed and closed depressions, and the West Okoboji Member for lake sediment associated with extant lakes.

**Source of name:** the De Forest Branch of Thompson Creek, Harrison County, Iowa, one of the watersheds originally studied by Daniels et al. (1963).

**Type Sections:** The original type sections were composed of loess-derived alluvium in a small western Iowa watershed (Daniels et al., 1963). Type sections for the Gunder and Roberts Creek members occur along Roberts Creek, Clayton County, in the Paleozoic Plateau region of northeastern Iowa. The type section for the Camp Creek Member occurs in Woodbury County in the Loess Hills of western Iowa, and the type section of the Corrington Member occurs in Cherokee County along the Little Sioux Valley in the Northwest Iowa Plains region.

**Description of Unit:** The DeForest Formation consists of fine-grained alluvium, colluvium, and pond sediments. A minor component of most members is sand or pebbly sand which, if present, is usually discontinuous, filling small scour channels at the base of the member or at the base of depositional units within members. Peat and muck occur in the Woden Member and infrequently as thin, local, discontinuous beds within the Gunder, Roberts Creek, and Camp Creek members.

Except where the tops of members have been erosionally truncated, soil profiles are developed in all members of the formation except the West Okoboji Member. Weakly expressed buried soils are locally preserved in all members except the Flack and West Okoboji. These buried soils reflect periods of landscape stability, but they are not widely traceable, even in individual drainage basins. They appear to record only short-lived local conditions. Secondary weathering-zone properties in the members vary with the depth and elevation of the water table.

**Nature of Contacts:** The DeForest Formation occurs at the land surface. It abruptly and unconformably overlies the Dows, Noah Creek, and any older Quaternary and Paleozoic formations into which it is incised. The contact is marked by an abrupt change in texture, sedimentary structures, and fossil content.

**Differentiation from other Units:** The alluvium, colluvium, and pond sediments of the DeForest Formation are generally unlike the deposits of any other formation on the DML. The Lake Mills Member of the Dows Formation consists of fine-grained sediment, but it tends to have a higher clay content and occurs in a different geomorphic setting (uplands instead of stream valleys). The Noah Creek Formation and the Pilot Knob Member of the Dows Formation are predominantly coarse sand and gravel. The Alden and Morgan members of the Dows Formation include poorly sorted diamicton deposits, which the DeForest Formation typically lacks. The Peoria Formation occurs on high terraces and uplands and is better sorted than DeForest formation deposits.

**Regional Extent and Thickness:** The DeForest Formation occurs in stream valleys, closed depressions, and on hillslopes across Iowa, and on the DML it also occurs in linked-depression drainageways. Thickness varies with geomorphic position and local relief. Where present, the formation varies in thickness from a few centimeters (inches) to several meters (greater than 20 feet) thick.

**Origin:** The DeForest Formation consists of post-glacial alluvium, colluvium, pond deposits, and organic sediment (peat and muck) that were deposited by or in water.

**Age:** The base of the DeForest Formation is time-transgressive. On the DML it is younger than 11,000 RCYBP in most areas, but is locally as old as 14,000 to 11,000 RCYBP. Deposition of the DeForest Formation continues to the present. The age of individual members is also time-transgressive, dependent on position in the drainage system and on geomorphic position.

### **Camp Creek Member**

**Source of Name:** Camp Creek a tributary of Garretson Drainage Ditch, Woodbury County, Iowa.

**Type Section:** Camp Creek cutbank exposure, Woodbury County, Iowa, NW 1/4, SW1/4 of section 1 T. 87 N., R. 45 W. (Bettis, 1990).

**Description of Unit:** Usually a calcareous to noncalcareous, very dark gray to brown, stratified (planar-bedded) silt loam to clay loam. Surface soils developed into the Camp Creek Member are Entisols (Typic Udifluvents). These soils consist of an organically enriched surface horizon (A horizon) grading to unaltered parent material. Where this unit is rapidly aggrading, surface soils are absent.

**Nature of Contacts:** This member is inset into or unconformably overlies the Gunder, Corrington and Roberts Creek members, depending on the local geomorphic setting and history of landuse. This unit often buries pre-settlement soil surface. May grade to sand and gravel in and adjacent to the modern channel belt.

**Differentiation from Other Members:** The Camp Creek Member differs from other members of the formation, in geomorphic position and nature of the stratigraphic sequence.

**Regional Extent and Thickness:** Thickness of the Camp Creek member is quite variable ranging from a few centimeters to over five meters (16.4 ft.).

**Origin:** Camp Creek Member consists of late-Holocene to post-settlement alluvium in and adjacent to modern channel belt, and at the base of steep slopes.

**Age:** Age is time-transgressive, dependent on drainage system and geomorphic position. In large valleys Camp Creek Member started aggrading as early as 400 B.P. and in small valleys as early as 150 B.P. It is still accumulating at present in both small and large valleys.

### **Roberts Creek Member**

**Source of Name:** Roberts Creek, Clayton County, Iowa

**Type Area:** Along Roberts Creek, Clayton County, Iowa, sections 6 and 7, T. 94 N., R. 5 W. (Baker, et al., 1996).

**Description of Unit:** Roberts Creek Member consists of dark, clayey, silty and loamy alluvium grading downward to sand and gravel; usually noneffervescent; thick sections are stratified at depth; detrital organic matter in lower part; relatively thick Mollisol (A-C or A-Bw-C profile) developed in the upper part (Bettis et al., 1992). Weakly expressed buried soils have been observed within the Roberts Creek Member, but these are not traceable from one valley to another. This unit includes the Mullenix and Turton members of Daniels, et al. (1963), which have been redesignated as beds within the Roberts Creek Member in the thick and moderately thick loess areas of western Iowa and adjacent states.

**Nature of Contacts:** Roberts Creek Member deposits overlie a wide variety of deposits including the Gunder and Corrington members, older alluvium, loess and glacial till.

**Differentiation From Other Members:** The Roberts Creek Member differs from other members of the formation, in geomorphic position and nature of the stratigraphic sequence. Soils are morphologically less well expressed and have darker B and C horizons than soils developed in the Gunder and Corrington members. The Roberts Creek Member is separated from younger DeForest Formation deposits (Camp Creek Member) by either a fluvial erosion surface or an unconformity marked by a buried soil (Bettis, 1995).

**Regional Extent and Thickness:** Roberts Creek deposits are found beneath flood plains of small and large valleys and often overlap Gunder Member deposits in 2<sup>nd</sup> and 3<sup>rd</sup>-order valleys. Unit thickness will vary dependent on size of valley. Usually unit thickness will vary from 1.5 to 5 m thick.

**Origin:** Roberts Creek Member consists of late-Holocene alluvium found in the modern floodplain, parallels the modern channel, also found in fan trenches.

**Age:** Unit age ranges from 4,000 to 500 B.P.

### **Gunder Member**

**Source of Name:** Roberts Creek, Clayton County, Iowa

**Type Area:** Along Roberts Creek, Clayton County, Iowa, sections 6 and 7, T. 94 N., R. 5 W. (Baker, et al., 1996).

**Description of Unit:** Gunder Member consists of oxidized brown to yellowish brown to grayish brown silt loam, silty clay loam, or loam grading to sand and gravel at depth. Usually noneffervescent, lower part may be stratified and reduced, detrital organic matter often present in lower coarse-grained part of unit; moderately well to somewhat poorly drained Mollisols and Alfisols developed in upper part. This member includes the Watkins and Hatcher members of Daniels et al. (1963) which have now been redesignated as beds within the Gunder Member. Buried soils are sometimes present within the Gunder Member, but are not traceable on a regional scale.

**Nature of Contacts:** Gunder Member deposits unconformably overlie loess, glacial till, bedrock, coarse alluvium, or organic-rich fine-grained alluvium. Overlying younger members of the formation are separated from the Gunder Member by a fluvial erosion surface or an unconformity marked by a buried soil.

**Differentiation From Other Members:** The Gunder Member differs from other members of the formation, in geomorphic position and nature of the stratigraphic sequence. Soils are morphologically better expressed and have lighter B and C horizons than soils developed in the Roberts Creek member.



**Regional Extent and Thickness:** Gunder deposits usually comprise low terrace that merges with side-slopes in a smooth concave upward profile. Usually unit thickness will vary from .5 to 4 m thick, with thickest deposits associated with Watkins member deposits.

**Origin:** Gunder Member consists of mid-early Holocene alluvium found low terrace positions merging with sideslopes.

**Age:** Unit ranges in age from 10,500 to 3000 B.P.

### Corrington Member

**Source of Name:** Corrington alluvial fan, Cherokee, County, Iowa

**Type Section:** Along the Little Sioux River Valley wall, Cherokee, County, Iowa, W 1/2, SW 1/4, SE 1/4 of section 4, T. 91 N., R. 40 W. (Hallberg, et al., 1974; Hoyer, 1980a, 1980b).

**Description of Unit:** The Corrington Member is the most internally variable unit of the formation and consists of very dark brown to yellowish brown oxidized loam to clay loam with interbedded lenses of sand and gravel; noneffervescent to effervescent at depth. The unit is stratified and usually contains several buried soils. Surface soils developed into this unit in are thick Mollisols (Cumlic Hapludolls) or Alfisols (Hapludalfs) that have argillic (Bt) horizons (Bettis, 1995).

**Nature of Contacts:** The Corrington Member buries coarse-grained older alluvium, glacial till, loess, or bedrock, and can grade laterally into Gunder Member deposits.

**Differentiation from Other Members:** The Camp Creek Member differs from other members of the formation, in geomorphic position and nature of the stratigraphic sequence. The presence of numerous buried soils (paleosols) and several fining-upward sequences often characterize unit.

**Regional Extent and Thickness:** Corrington Member deposits compose alluvial fans located where small and moderate-size valleys (2<sup>nd</sup>-and 3<sup>rd</sup>-order) enter larger valleys. Fans will vary in thickness, typically thicker sections have been measured in western Iowa. At the type section in Cherokee County, section thickness was measured at 11 m (36 ft.).

**Origin:** Corrington Member is found in alluvial fans and colluvial slopes along the margins of large to moderate-size valley. Deposits are variably textured and accumulated by channeled flow, sheetwash, and debris flow (Hoyer, 1980b).

**Age:** Unit ranges in age from 9000 to about 2500 B.P

## PEORIA FORMATION

The Peoria Formation consists of wind-transported sediments and occurs throughout Iowa.

**Source of name:** the city of Peoria, Peoria County, Illinois.

**Type Section:** the Tindall School Section, a borrow pit in the west bluff of the Illinois Valley south of Peoria, Peoria County, Illinois, in the SW 1/4, SW 1/4, NE 1/4 of section 31, T. 7 N., R. 6 E. (Willman and Frye, 1970).

**Description of Unit:** The Peoria Formation includes wind-transported sediments. Two facies are recognized in Iowa, a silt facies (loess) and a sand facies (eolian sand). The sediments are well sorted and the two facies may be interbedded. Textures range from silt loam to medium-to-fine sand. Macroscopic bedding structures are rare and are found primarily in locations proximal to a valley source where the formation's sediments are thick. Where present, bedding structures include planar beds with inverse grading in the silt facies, and planar beds to steep foresets in the sand facies. Where eolian sand overlies sand-and-gravel deposits of the Noah Creek Formation it is included in that formation. On the DML, secondary pedogenic alteration has modified most Peoria Formation deposits.

**Nature of Contacts:** The Peoria Formation usually occurs at the land surface. It abruptly and unconformably overlies older Quaternary formations and paleosols developed in them. Beneath the DML the silt facies of the formation is buried by Dows Formation glacial diamicton, while the sand facies occurs at the land surface and abruptly and unconformably overlies the Dows Formation. The contact with other

units is marked by an abrupt change in texture, sedimentary structures, fossil content, or secondary weathering characteristics.

**Differentiation From Other Units:** The wind-sorted sediments of the Peoria Formation are generally unlike the deposits of any other formation on the DML. The Lake Mills Member of the Dows Formation consists of fine-grained sediment, but it has greater variability, a higher clay content, and occurs in a different geomorphic setting. The Noah Creek Formation and Pilot Knob Member of the Dows Formation are more poorly sorted and contain coarse sand and gravel. The DeForest Formation contains some sandy sediment, but the bedding structures and sorting of these are distinct from those associated with the Peoria Formation.

**Regional Extent and Thickness:** The Peoria Formation occurs on uplands and high terraces throughout Iowa. In north-central Iowa, the silt facies of the formation is buried by glacial diamicton of the Dows Formation, except in very restricted, small areas adjacent to major river valleys in the southern part of the lobe. On the DML, the formation is usually restricted to a narrow belt on the upland along major stream valleys. Thickness varies with respect to distance from the valley source. Proximal to the Missouri Valley in western Iowa, the formation usually is more than thirty meters (90 ft) thick. On the DML the formation ranges from a few centimeters to about three meters (9 ft) in thickness.

**Origin:** The Peoria Formation consists of wind-deposited sediment. The formation's sediments were derived from wind reworking of valley-train outwash. The sand facies also includes sediments reworked from older eolian sand deposits.

**Age:** The Peoria Formation is time transgressive. The silt facies was deposited between about 22,000 and 12,500 RCYBP, while the sand facies includes deposits that accumulated contemporaneous with the silt facies, as well as others that accumulated during the Holocene to the present. Most Peoria Formation deposits on the DML accumulated between about 14,000 and 11,000 RCYBP, and have undergone various degrees of wind reworking during the Holocene.

## PISGAH FORMATION

The Pisgah Formation includes both primary eolian silt and eolian silt and related slope sediments that have been altered by a combination of colluvial hillslope processes, pedogenic and periglacial processes.

**Source of name:** the town of Pisgah, Harrison County, Iowa.

**Type Section:** the Loveland Paratype Section, cut north of Interstate 680 at intersection with Interstate 29, Pottawattamie County, Iowa, in the NW1/4, SE 1/4, Sec. 3 T77N R44W

**Description of Unit:** The Pisgah Formation is primary eolian silt (loess) and eolian silt and related hillslope sediments that have been modified by a combination of colluvial hillslope processes, pedogenic and periglacial processes. As a result of various mixing processes most Pisgah Formation deposits have a coarse, gritty silt loam texture. Throughout the state, the unit is usually thin (0.15 to 1 m thick), and has been pedogenically altered both at the base, by slow incorporation into the existing Sangamon Geosol, and at the top, from pedogenesis producing the Farmdale Geosol before burial by the Peoria Formation (Kemmis et al., 1992). Typically, the Farmdale Geosol is expressed as a thin, dark grayish brown, weakly developed buried soil consisting of a leached A-C or E-C profile with platy or granular structure and associated charcoal flecks. Where the Pisgah Formation is thin (<30cm) the Farmdale Geosol is developed through the entire unit.

**Nature of Contacts:** The Pisgah Formation is typically buried by the Peoria Formation (silt facies--loess). Beneath the Des Moines Lobe the formation is buried Peoria loess, which in turn is buried by Dows Formation glacial diamicton. A prominent feature at the Peoria/ Pisgah contact is an involuted contact which is attributed to Wisconsin-age periglacial activity. Pisgah deposits typically overlie the Sangamon Geosol that is developed in Loveland or Pre-Wisconsin-age loess(s) or undifferentiated Pre-Illinoian glacial till.

**Differentiation From Other Units:** The wind-transported and slope reworked sediments of the Pisgah Formation are unlike the deposits of the Dows and Noah Creek formations on the DML. The Lake Mills Member of the Dows Formation consists of fine-grained silty sediment, but it has greater variability, a

higher clay content, and occurs in a different geomorphic setting. The Noah Creek Formation and Pilot Knob Member of the Dows Formation are more poorly sorted and contain coarse sand and gravel. The DeForest Formation contains some silty sediment, but the bedding structures and sorting of these are distinct from those associated with the Pisgah Formation, and occurs in a different geomorphic setting. The Pisgah Formation is genetically similar to the Peoria Formation, however it has been noticeably altered by pedogenic, colluvial hillslope and periglacial processes. The Pisgah Formation was referred to as the “basal Wisconsin loess” in previous literature (Ruhe, 1969).

**Regional Extent and Thickness:** The Pisgah Formation is preserved on relatively stable upland surfaces, but is absent on Iowan Erosion Surface of northeastern Iowa and on “Iowan” steps of the stepped erosion surface landscapes associated with the Southern Drift Plain landform region of the state (Hallberg et al., 1978). Where present on the DML, the formation is buried by Dows Formation glacial till and Peoria Formation loess. Thickness varies with landscape position and distance from the eolian source. In central and eastern Iowa, the formation is commonly only 0.15 to 1 m thick. In extreme western Iowa, adjacent to the Missouri River valley the unit is 3 to 4 m thick. The Pisgah Formation stratigraphic position is equivalent to the Roxana Silt of Illinois and the Gilman Canyon Formation of Nebraska, but lithologic properties are different (Bettis, 1990)

**Origin:** The Pisgah Formation originated as eolian silt that has been altered by a combination of colluvial hillslope processes, pedogenic and periglacial processes.

**Age:** The Pisgah Formation is time transgressive. The wind-transported silt was deposited between about 45,000 and 25,500 RCYBP while Peoria loess began to accumulate shortly after 25,000 RCYBP (Bettis, 1990). Dates from the Farmdale Geosol, developed in the upper part of the Pisgah Formation indicate that ages range from 28,000 to 16,500 RCYBP with ages decreasing with distance from the Missouri River valley (Ruhe, 1976).

## GLASFORD FORMATION

Although several Glasford Formation members are recognized in Illinois, the only Glasford Formation member present in Iowa is the Kellerville Member. The Glasford Formation is the most widespread glacial unit in Illinois, but only reaches into Iowa in the southeastern corner.

**Source of name:** Glasford, Peoria County, Illinois.

**Type Section:** the type section for the Glasford Formation is located in Peoria County at the Tindall School Section, SW SW NE S31, T7N, R6E (Willman and Frye, 1970).

**Description of Unit:** In Iowa, the Glasford Formation Kellerville Member is separated into two till facies (a subglacial or basal till facies and a superglacial facies) based on stratigraphic position, sedimentological properties, and the consistency-density-consolidation properties (Hallberg et al., 1980; Wickham, 1980; Lineback, 1979). The two facies are similar in mineralogy (both clays and sand-fraction lithology), but not physical characteristics. The subglacial till facies has a firm, dense uniform till matrix with texture ranging from silt loam to a light clay loam (on average a loam till relatively high in silt). In contrast, the superglacial facies may be composed of till, diamicton (reworked till such as superglacial debris flows), sorted fluvial and lacustrine sediments, and peat beds. Deposits may be interbedded or occur as a contorted melange of sediments. The superglacial till is highly variable in density and texture, and contains a wide variety of sediments. The texture values tend to cluster if samples are from the same general area. The overall range of matrix texture includes sandy loam, loam, silt loam, silty clay loam, clay loam, and clay. The stratified deposits within the superglacial till facies vary in texture from sand, sand and gravel, to very heavy clay. The superglacial facies also contains some peats and organic silts (Hallberg and Baker, 1980). Although the superglacial and subglacial facies vary widely in texture, they are very similar in mineralogy, especially the clay abundance and the sand-fraction lithology. The clays typically contain 46% expandables, 34% illite, and 20% kaolinite plus chlorite, but may grade vertically and laterally.

**Nature of Contacts:** The Glasford Formation is underlain by the Petersburg Silt, or where absent the Yarmouth Soil (Willman and Frye, 1970). The Yarmouth Soil may be formed in Pre-Illinoian till, allu-

vium or local bedrock. The Sangamon Soil is typically formed in the upper surface of the Glasford Formation, but in areas where extensive erosion has occurred it may be directly overlain by loess (Pisgah or Peoria formation) or alluvium.

**Differentiation from other Units:** Clay mineral characteristics (including a high illite to kaolinite plus chlorite ratio, moderate amounts of expandables, and the frequent occurrence of identifiable chlorite peaks) are used to distinguish the Illinoian age tills from other units in the area. Clay mineral data from the Glasford Formation are more variable (wider range) than for the Pre-Illinoian tills (Hallberg et al., 1980; Lineback, 1979). The Kellerville Member of the Glasford Formation is distinguished from the Wolf Creek Formation by the relatively high illite and dolomite contents. Additionally, the Kellerville Member exhibits an abundance of Pennsylvanian lithologies in the very coarse sand through cobble size particles (Hallberg, 1980b).

**Regional Extent and Thickness:** The Lake Michigan Lobe of Illinoian ice did not reach far into Iowa, only advancing into the most southeastern portion of the state. Leverett (1898) originally defined the terminal boundary, and his distribution map in Iowa has only been slightly modified since (Kay and Graham, 1943; Ruhe, 1969). The Illinoian deposits in southeast Iowa extend along the western edge of the Mississippi River from just south of Fort Madison northward to near the mouth of the Wapsipinicon River at the boundary between Scott and Clinton Counties (Leverett, 1898, Kay and Graham, 1943; Ruhe, 1969). The distance west from the Mississippi River varies, ranging from approximately 6.5 km (4 miles) near Muscatine to 32 km (20 miles) north of Burlington. Illinoian deposits have been identified in Lee (NE ¼), Des Moines (all but NW corner), Henry (only SE corner), Louisa (eastern 2/3), Muscatine (all but NW ¼), and Scott Counties (southern 2/3). The superglacial facies varies in thickness from a few feet to 93 feet (28.4m) at the terminal ridge of the Kellerville at Yarmouth (Hallberg et al., 1980). The thickest known section of Kellerville Till in Iowa is located at the Yarmouth core site near the western terminus; however, the Illinoian materials in the core are entirely made up of the superglacial facies (Hallberg et al., 1980). Within a few miles of the Mississippi River at the Nelson Quarry Section, 77 feet (23.5m) of the basal till facies is present (Hallberg et al., 1980). Therefore, in areas where both the superglacial and subglacial till facies are present, the total thickness could exceed 93 feet. The thickness throughout most of western Illinois ranges from less than 1m to 25m thick, but estimates indicate that it may be up to 50m thick in deep bedrock valleys (Wickham, 1980).

**Origin:** The Glasford Formation consists of glacial tills and intercalated outwash deposits associated with the advance of the Lake Michigan Lobe from the northeast.

### **Kellerville Member**

The Kellerville Till Member is the only Glasford Formation till present in Iowa. Willman and Frye (1970) used the term to replace the Mendon Till (Frye et al., 1964 and 1969) and Payson Till (Leighton and Willman, 1950; Wanless, 1957).

**Source of name:** Kellerville, Adams County, Illinois

**Type Section:** the type section for the Kellerville Till Member is defined from two roadcut exposures 2 miles southwest of Kellerville in the Washington Grove School Section- NW NW SW S11, T2S, R5W (Willman and Frye, 1970). The type section is approximately 100km south of Burlington, Iowa.

**Description of Unit:** The Kellerville Till Member is separated into two till facies (a subglacial or basal till facies and a superglacial facies) based on stratigraphic position, sedimentological properties, and the consistency-density-consolidation properties (Hallberg et al., 1980; Wickham, 1980; Lineback, 1979). The two facies are similar in mineralogy (both clays and sand-fraction lithology), but not physical characteristics. The subglacial till facies has a firm, dense uniform till matrix with texture ranging from silt loam to a light clay loam (on average a loam till relatively high in silt). In contrast, the superglacial facies may be composed of till, diamicton (reworked till such as superglacial debris flows), sorted fluvial and lacustrine sediments, and peat beds. Deposits may be interbedded or occur as a contorted melange of sediments. The superglacial till is highly variable in density and texture, and contains a wide variety of sediments. The texture values tend to cluster if samples are from the same general area. The overall

range of matrix texture includes sandy loam, loam, silt loam, silty clay loam, clay loam, and clay. The stratified deposits within the superglacial till facies vary in texture from sand, sand and gravel, to very heavy clay. The superglacial facies also contains some peats and organic silts (Hallberg and Baker, 1980). Although the superglacial and subglacial facies vary widely in texture, they are very similar in mineralogy, especially the clay abundance and the sand-fraction lithology. The clays typically contain 46% expandables, 34% illite, and 20% kaolinite plus chlorite.

**Nature of Contacts:** The Kellerville Till Member is underlain by the Petersburg Silt, or where absent the Yarmouth Soil (Willman and Frye, 1970). The Yarmouth Soil may be formed in Pre-Illinoian till, alluvium or local bedrock. The Sangamon Soil is typically formed in the upper surface of the Kellerville, but in areas where extensive erosion has occurred it may be directly overlain by loess (Pisgah or Peoria formation) or alluvium.

**Differentiation from other Units:** Additional clay mineral characteristics (including a high illite to kaolinite plus chlorite ratio, moderate amounts of expandables, and the frequent occurrence of identifiable chlorite peaks) are used to distinguish the Kellerville from other units in the area. Clay mineral data from the Kellerville Member are more variable (wider range) than for the Pre-Illinoian tills (Hallberg et al., 1980; Lineback, 1979). The Kellerville Member of the Glasford Formation is distinguished from the Wolf Creek Formation by the relatively high illite and dolomite contents. Additionally, the Kellerville Member exhibits an abundance of Pennsylvanian lithologies in the very coarse sand through cobble size particles (Hallberg, 1980b). The Wolf Creek Formation also has a much higher limestone to dolomite ratio (greater than 0.40) than the Kellerville Formation deposits.

**Regional Extent and Thickness:** The Lake Michigan Lobe of Illinoian ice did not reach far into Iowa, only advancing into the most southeastern portion of the state. Leverett (1898) originally defined the terminal boundary, and his distribution map in Iowa has only been slightly modified since (Kay and Graham, 1943; Ruhe, 1969). The Illinoian deposits in southeast Iowa extend along the western edge of the Mississippi River from just south of Fort Madison northward to near the mouth of the Wapsipicon River at the boundary between Scott and Clinton Counties (Leverett, 1898, Kay and Graham, 1943; Ruhe, 1969). The distance west from the Mississippi River varies, ranging from approximately 6.5 km (4 miles) near Muscatine to 32 km (20 miles) north of Burlington. Illinoian deposits have been identified in Lee (NE ¼), Des Moines (all but NW corner), Henry (only SE corner), Louisa (eastern 2/3), Muscatine (all but NW ¼), and Scott Counties (southern 2/3). The superglacial facies varies in thickness from a few feet to 93 feet (28.4m) at the terminal ridge of the Kellerville at Yarmouth (Hallberg et al., 1980). The thickest known section of Kellerville Till in Iowa is located at the Yarmouth core site near the western terminus; however, the Illinoian materials in the core are entirely made up of the superglacial facies (Hallberg et al., 1980). Within a few miles of the Mississippi River at the Nelson Quarry Section, 77 feet (23.5m) of the basal till facies is present (Hallberg et al., 1980). Therefore, in areas where both the superglacial and subglacial till facies are present, the total thickness could exceed 93 feet. The thickness throughout most of western Illinois ranges from less than 1m to 25m thick, but estimates indicate that it may be up to 50m thick in deep bedrock valleys (Wickham, 1980).

**Origin:** Studies have determined that the Illinoian till was deposited by the Lake Michigan Lobe which advanced across western Illinois from the northeast and into Iowa (Leverett, 1898; Wickham, 1980). The Lake Michigan Lobe incorporated Paleozoic bedrock from the Lake Michigan Basin, which is reflected in both the mineralogy of the matrix and clasts of the Illinoian deposits (Lineback, 1980; Wickham, 1980). The northeastern source area for the Illinoian age tills is also evidenced by till fabrics and glacial landform orientations (Lineback, 1979).

## WOLF CREEK FORMATION

The Wolf Creek Formation is subdivided into three members (oldest to youngest): the Winthrop, Aurora and Hickory Hills members. Information on the formation as a whole is presented first, followed by more specific descriptions for individual members.

**Source of name:** Wolf Creek, northern Tama County, Iowa.

**Type Section:** the type area for the Wolf Creek Formation is defined from several reference localities in the region around Wolf Creek in Geneseo, Clark, Buckingham, and Grant Townships, in northern Tama County.

**Description of Unit:** The Wolf Creek Formation is predominantly a massive, uniform, basal till, but may also include fluvial silts, sands and gravels and local fine-textured swale fill deposits and peat. On average the texture is loam, but subtle differences may be used to help distinguish members. The Wolf Creek Formation averages 50-60% expandable clays (slightly lower in the southeastern portion of the state), 16-19% illite, and 22-24% kaolinite plus chlorite (Hallberg et al., 1980).

**Nature of Contacts:** In areas of southeast Iowa that were glaciated during the Illinoian, the upper boundary of the Wolf Creek Formation is marked by the unconformable contact with deposits of the Glasford Formation. Where Illinoian age deposits are present, the Yarmouth Paleosol is formed in the top of the Wolf Creek Formation. Beyond the reaches of the Illinoian deposits, the Yarmouth-Sangamon Paleosol is developed in the Wolf Creek. The individual till members may be directly overlain by each other or be separated by undifferentiated sediments, glaciofluvial deposits, or paleosols. The Wolf Creek Formation is underlain by either the Alburnett Formation or Paleozoic bedrock.

**Differentiation from other Units:** The Wolf Creek Formation is distinguished from the Kellerville Member of the Glasford Formation (Illinoian age till) by the relatively low illite and dolomite contents. Additionally, the Kellerville Member exhibits an abundance of Pennsylvanian lithologies in the very coarse sand through cobble size particles (Hallberg 1980b). The Wolf Creek Formation also has a much higher limestone to dolomite ratio (greater than 0.40) than the Kellerville Formation deposits.

Differentiation of the Wolf Creek and Alburnett formation tills is difficult in the field without the assistance of stratigraphic boundaries, but mineralogical characteristics can be used to distinguish them. The most useful characteristic to differentiate between the Wolf Creek and Alburnett formations is the clay mineralogy. The Wolf Creek Formation has a higher expandable clay percentage, averaging around 62%; whereas the Alburnett Formation has lower expandable clay percentages, near 43% (Hallberg et al., 1980a).

**Origin:** The Wolf Creek Formation consists of three till members associated with several Pre-Illinoian ice advances. Based on the physical properties of the majority of the Wolf Creek Formation deposits (massive structure, high density, uniform texture) they likely represent a basal, or subglacial, till facies.

**Age and Correlation:** The Wolf Creek Formation represents the youngest of the Pre-Illinoian glaciations. Pre-Illinoian deposits in Iowa range from older than 2.2 million years to approximately 500,000 years ago based on volcanic ash dates in western Iowa (Easterbrook and Boellstorff, 1984; Hallberg, 1986). Paleomagnetic studies of the Wolf Creek Formation in east-central Iowa indicate that these deposits have normal polarity and are therefore younger than 790,000 years (Baker and Stewart, 1984).

### Winthrop Member

**Source of name:** the town of Winthrop, Buchanan County, Iowa.

**Type Section:** the type area of the Winthrop Till Member consists of a railroad cut and drill-core section approximately 1¼ miles (2 km) west of Winthrop in Buchanan County in the NW¼ of Section 3, Township 88N, Range 8W. The exposure is about 5 ½ miles (8.8 km) east of Independence and 2 ¼ miles (3.6 km) east of Doris Station.

**Description of Unit:** The Winthrop Till Member is the oldest and least well-known of the Wolf Creek Formation tills due to poor preservation. The color varies within the weathering profile from a light-yellowish brown to dark gray. Texturally, the Winthrop Till Member is a loam to light clay loam with averages of 25% clay, 41% silt and 34% sand. Generally, it contains more silt than sand. Clay mineral percentages in the Winthrop Till Member average 60% ±4.3 (range 51-68) expandables, 17% ±2.2 (range 10-20) illite, and 24% ±3.8 (range 16-31) kaolinite plus chlorite (Hallberg, 1980a). The Winthrop Till Member tends to have slightly higher values of kaolinite than the other tills of the Wolf Creek Formation. The Winthrop Till Member exhibits a high limestone to dolomite ratio (median and mode >15) and dolomite is commonly absent.

**Nature of Contacts:** The Winthrop Till Member is overlain by either leached unnamed sediments separating it from the Aurora Till Member, the Aurora Till Member, younger Wolf Creek Formation deposits, or may be exhumed as the surface till where erosion was severe enough to remove the younger deposits.

The lower boundary of the Winthrop Till Member is equally complex, with the contact being marked by the underlying bedrock, sediments of the Alburnett Formation, or in the most complete sections by the top of the Westburg Paleosol. Where present, the Westburg Paleosol occurs below the Winthrop Till Member of the Wolf Creek Formation and is developed in deposits of the Alburnett Formation or older rock units.

**Differentiation from other Units:** The Winthrop Member is distinguished from the Kellerville Member of the Glasford Formation (Illinoian age till) by the relatively low illite and dolomite contents. The Kellerville also exhibits an abundance of Pennsylvanian lithologies in the very coarse sand through cobble size particles and a much lower limestone to dolomite ratio (less than 0.40) (Hallberg, 1980b). All members of the Wolf Creek Formation have average limestone to dolomite ratios greater than 0.40.

Differentiation of the Wolf Creek Formation members and Alburnett Formation materials is difficult in the field without the assistance of stratigraphic boundaries, but mineralogical characteristics can be used to distinguish them. The most useful characteristic to distinguish between the Wolf Creek members and the Alburnett Formation is the clay mineralogy (Hallberg, 1980a). The Wolf Creek Formation has higher expandable clay percentages, averaging around 62%; whereas the Alburnett Formation has lower expandable clay percentages, near 43%.

Clay mineral variation cannot be utilized to differentiate members of the Wolf Creek Formation, and the sand-fraction lithology generally overlaps. However, the limestone to dolomite ratio and grain-size distribution has been useful for discriminating between members (Hallberg, 1980a). Differentiation can be difficult in areas of isolated exposures where only one till is exposed (or the till varies to an end member within its range). Typically, the Hickory Hills member almost always has more sand than silt and has higher values for total carbonate and sedimentary grains than the Aurora or Winthrop members. The Hickory Hills member also has a low limestone to dolomite ratio. Both the Aurora and Winthrop members have more silt than sand, a high limestone to dolomite ratio, and often do not have dolomite. Overall, these similarities between the Aurora and Winthrop members make it difficult to distinguish the two if only one is present. The Winthrop generally has lower values for total carbonates and sedimentary grains.

**Regional Extent and Thickness:** Due to the limited number of positive identifications, the thickness of the Winthrop Till Member is poorly known and difficult to determine. In the composite Winthrop locality it varies from 2 to about 15 feet (0.6-4.6 m), and has a thickness of 48 feet (14.6 m) in the 4-Mile Creek area.

**Origin:** The Winthrop Member of the Wolf Creek Formation consists of deposits associated with an advance of Pre-Illinoian ice. Based on the physical properties of these deposits (massive structure, high density, uniform texture) they likely represent a basal, or subglacial, till facies.

**Age and Correlation:** The Wolf Creek Formation members represent the youngest of the Pre-Illinoian glaciations. Pre-Illinoian deposits in Iowa range from older than 2.2 million years to approximately 500,000 years ago based on volcanic ash dates in western Iowa (Easterbrook and Boellstorff, 1984; Hallberg, 1986). Paleomagnetic studies of the Wolf Creek Formation in east-central Iowa indicate that these deposits have normal polarity and are therefore younger than 790,000 years (Baker and Stewart, 1984).

### **Aurora Member**

**Source of name:** the town of Aurora, Buchanan County, Iowa.

**Type Section:** the type area for the Aurora Till Member is the Aurora Transect located approximately 2 miles (3.2 km) southwest of the town of Aurora in northeast Buchanan County, Iowa. The transect consists of core-holes drilled along the axis of the stepped erosion surfaces in the area of the regional divide between the Wapsipinicon and Maquoketa Rivers in the NE1/4 of Section 23, T 90N, R 8W.

**Description of Unit:** The Aurora Till Member is a basal till with relatively uniform characteristics. The texture of the Aurora Till Member is loam, averaging 22% clay, 40% silt, and 38% sand. The Aurora has a high limestone to dolomite ratio (median and mode >15), and often no dolomite is present. Color varies

vertically within the weathering profile from light yellowish-brown to dark gray or dark greenish gray. Clay mineralogy averages 62 %  $\pm$ 3.6 (range 55-70) expandables, 18%  $\pm$ 2.5 (range 13-24) illite, and 21%  $\pm$ 2.3 (range 17-24) kaolinite plus chlorite (Hallberg, 1980a).

**Nature of Contacts:** The upper boundary of the Aurora Till Member varies in relation to the amount of erosion. In complete sections, this boundary is marked by the contact with the unnamed weathered (leached) sediments, which generally contain the Dysart Paleosol (which is overlain by the Hickory Hills Till Member). Where the Dysart Paleosol has been eroded, the contact may be directly with the Hickory Hills Till Member or a sharp diffuse contact zone including glaciofluvial sediments of the Hickory Hills Till Member. In some places erosion is severe enough that the Aurora Till Member is the surficial till unit and may be overlain by a thin veneer of Wisconsin to Holocene surficial sediments or eolian sand, Wisconsinan loess, or Pre-Wisconsinan sediments.

In complete sections, the lower boundary of the Aurora Till Member may be marked by the contact with leached sediments and weak paleosols separating it from the Winthrop Till Member. In areas where erosion has occurred, the Aurora Member may be in direct contact with the Winthrop Till Member, various sediments of the Alburnett Formation, or bedrock. If either of the last two settings is the case, the Aurora Till Member also marks the base of the Wolf Creek Formation.

**Differentiation from other Units:** The Wolf Creek Formation tills are distinguished from the Kellerville Member of the Glasford Formation (Illinoian age till) by the relatively low illite and dolomite contents. The Kellerville also exhibits an abundance of Pennsylvanian lithologies in the very coarse sand through cobble size particles (Hallberg 1980b). The Kellerville has a much lower limestone to dolomite ratio (less than 0.40) than the Wolf Creek Formation deposits (95% of which all have limestone to dolomite ratios greater than 0.40).

Differentiation of the Wolf Creek Formation members and Alburnett Formation materials is difficult in the field without the assistance of stratigraphic boundaries, but mineralogical characteristics can be used to distinguish them. The most useful characteristic to distinguish between the Wolf Creek and Alburnett Formations are the clay mineral percentages. The Wolf Creek Formation has higher expandable clay values, averaging around 62%; whereas the Alburnett Formation has lower expandable clay percentages, near 43% (Hallberg, 1980a).

Clay mineral variation cannot be utilized to differentiate members of the Wolf Creek, and the sand-fraction lithology generally overlaps. However, the limestone to dolomite ratio and grain-size distribution has been useful for discriminating between members (Hallberg, 1980a). Differentiation can be difficult in areas of isolated exposures where only one till is exposed (or the till varies to an end member within its range). Typically, the Hickory Hills member almost always has more sand than silt and has higher values for total carbonate and sedimentary grains than the Aurora or Winthrop members. The Hickory Hills member also has a low limestone to dolomite ratio. Both the Aurora and Winthrop members have more silt than sand, a high limestone to dolomite ratio, and often do not have dolomite. Overall, these similarities between the Aurora and Winthrop members make it difficult to distinguish the two if only one is present. The Winthrop generally has lower values for total carbonates and sedimentary grains.

**Regional Extent and Thickness:** The thickness of the Aurora Member is highly variable. In some areas it has been entirely removed by erosion, and at the 4-Mile Creek locality it may reach 100 feet (31m) in thickness. In most areas it ranges from 20 to 35 feet (6 to 11 m) in thickness.

**Origin:** The Aurora Member of the Wolf Creek Formation consists of deposits associated with an advance of Pre-Illinoian ice. Based on the physical properties of these deposits (massive structure, high density, uniform texture) they likely represent a basal, or subglacial, till facies.

**Age and Correlation:** The Wolf Creek Formation members represent the youngest of the Pre-Illinoian glaciations. Pre-Illinoian deposits in Iowa range from older than 2.2 million years to approximately 500,000 years ago based on volcanic ash dates in western Iowa (Easterbrook and Boellstorff, 1984; Hallberg, 1986). Paleomagnetic studies of the Wolf Creek Formation in east-central Iowa indicate that these deposits have normal polarity and are therefore younger than 790,000 years (Baker and Stewart, 1984).



## Hickory Hills Member

**Source of name:** Hickory Hills Park, NW1/4 of the SE1/4 of section 10, T 86N, R 13W (Geneseo Township), Tama County.

**Type Section:** the type area of the Hickory Hills Till Member is within the vicinity of Hickory Hills Park. Two principal reference localities are described within the type area. The 402 Road cut Section is designated the type locality and section. Casey's Paha East is the principal reference locality. Several other reference localities (the Hayward's Paha Transect and Buckingham Section) are needed to fully describe the upper and lower boundaries.

**Description of Unit:** Due to weathering, the Hickory Hills Till Member varies vertically in color from light-yellowish brown (10YR 5/6-8) in the oxidized zone to dark greenish gray (5GY 4/1) in the unoxidized zone. Texturally, the Hickory Hills Till Member is a loam, averaging about 22% clay, 34% silt, and 44% sand (Hallberg, 1980a). In thick sections, the till tends to be quite uniform texturally, but where it is thin it tends toward a mixed composition incorporating the material below. The Hickory Hills Member almost always has more sand than silt. The average clay mineral percentages are 63%  $\pm$ 4.5 (range 52-73) expandables, 17%  $\pm$ 3.3 (range 11-23) illite, and 20%  $\pm$ 2.2 (range 14-25) kaolinite plus chlorite (Hallberg, 1980a). The Hickory Hills till has a lower limestone to dolomite ratio than the other Wolf Creek Formation members.

**Nature of Contacts:** The lower boundary is commonly marked by the contact with the Dysart Paleosol and related unnamed sediments, or where absent it rests directly on the Aurora Till Member. When resting directly on the Aurora, the boundary is often a complex zone of contorted glaciofluvial sediments related to the Hickory Hills Till Member. If pre-Hickory Hills Till erosion was extensive, the Hickory Hills Till Member may lie directly on any older unit from the Winthrop Till Member to Paleozoic bedrock. In some sections the contact with the Dysart Paleosol is not clear due to block inclusions of the Dysart Paleosol that were sheared into the lower portion of the Hickory Hills Till Member. The upper boundary is also complex due to erosion, and it may be overlain by the Yarmouth- Sangamon or Late Sangamon surface, Wisconsinan age sediments or other surficial materials.

**Differentiation from other Units:** The Wolf Creek Formation tills are distinguished from the Kellerville member of the Glasford Formation (Illinoian age till) by the relatively low illite and dolomite contents. The Kellerville also exhibits an abundance of Pennsylvanian lithologies in the very coarse sand through cobble size particles (Hallberg 1980b). The Kellerville has a much lower limestone to dolomite ratio (less than 0.40) than the Wolf Creek Formation tills (95% of which all have limestone to dolomite ratios greater than 0.40).

Differentiation of the Wolf Creek and Alburnett Formation tills is difficult in the field without the assistance of stratigraphic boundaries, but mineralogical characteristics can be used to distinguish them. The most useful characteristic to distinguish between the Wolf Creek and Alburnett Formations is the clay mineralogy (Hallberg, 1980a). The Wolf Creek Formation has higher expandable clay percentages, averaging around 62%; whereas the Alburnett Formation has lower expandable clay percentages, near 43%.

Clay mineral variation cannot be utilized to differentiate members of the Wolf Creek Formation, and the sand-fraction lithology generally overlaps. However, the limestone to dolomite ratio and grain-size distribution has been useful for discriminating between members (Hallberg, 1980a). Differentiation can be difficult in areas of isolated exposures where only one till is exposed (or the till varies to an end member within its range). Typically, the Hickory Hills member almost always has more sand than silt and has higher values for total carbonate and sedimentary grains than the Aurora or Winthrop members. The Hickory Hills member also has a low limestone to dolomite ratio. Both the Aurora and Winthrop members have more silt than sand, a high limestone to dolomite ratio, and often do not have dolomite. Overall, these similarities between the Aurora and Winthrop members make it difficult to distinguish the two if only one is present. The Winthrop generally has lower values for total carbonates and sedimentary grains.

**Regional Extent and Thickness:** The thickness of the Hickory Hills Till Member is extremely variable. Due to erosion it is absent in some areas, however, in more complete sections the member ranges from 10 to over 50ft (3-15m).

**Origin:** The Hickory Hills Member of the Wolf Creek Formation consists of deposits associated with an advance of Pre-Illinoian ice. Based on the physical properties of these deposits (massive structure, high density, uniform texture) they likely represent a basal, or subglacial, till facies.

**Age and Correlation:** The Wolf Creek Formation members represent the youngest of the Pre-Illinoian glaciations. Pre-Illinoian deposits in Iowa range from older than 2.2 million years to approximately 500,000 years ago based on volcanic ash dates in western Iowa (Easterbrook and Boellstorff, 1984; Hallberg, 1986). Paleomagnetic studies of the Wolf Creek Formation in east-central Iowa indicate that these deposits have normal polarity and are therefore younger than 790,000 years (Baker and Stewart, 1984).

## ALBURNETT FORMATION

The Alburnett Formation is separated into several “undifferentiated” members. No consistent discriminating characteristics are available for these members, and the only differentiation comes where stratigraphic position allows. Therefore, the following description will be used for all the members.

**Source of name:** the town of Alburnett, Linn County, Iowa.

**Type Section:** the region around the town of Alburnett, Otter Creek Township (T 85N, R 7W), Linn County, Iowa.

**Description of Unit:** The Alburnett Formation is composed of multiple till units, which are "undifferentiated", and a variety of fluvial deposits. Minor paleosols may also be identified within the deposits. Throughout eastern Iowa, these deposits fill and bury the deep bedrock channels. The till is typically a uniform, massive, basal till. The Alburnett Formation ranges in color from light-yellowish brown in the oxidized zone, to dark gray or dark-greenish gray in the unoxidized zone.

The Alburnett Formation is defined by its stratigraphic position and distinctive clay mineralogy. The tills are generally loam textured, but range to light clay loam. On average, the Alburnett Formation consists of 18.7% clay, 36.8% silt, and 44.4% sand. The Alburnett Formation contains 44% expandables, 24% illite, and 32% kaolinite plus chlorite. In comparison with the Wolf Creek Formation, the Alburnett tills have significantly lower percentages of expandable clay minerals and higher kaolinite plus chlorite (Hallberg et al., 1980).

**Nature of Contacts:** The upper boundary of the Alburnett Formation is an unconformity of variable magnitude. Where the section is complete, the top of the Westburg Paleosol marks the upper boundary and is overlain by the Winthrop Till Member of the Wolf Creek Formation. Where the paleosol is eroded, any member of the Wolf Creek Formation, Wisconsin loess or other surficial sediments may overlie the Alburnett Formation. The lower boundary of the Alburnett Formation is marked by an unconformable contact with the bedrock. Glaciofluvial deposits may also be located at the base of the Alburnett Formation.

**Differentiation from other Units:** Pre-Illinoian tills are distinguished from the Kellerville Member of the Glasford Formation (Illinoian age till) by the relatively low illite and dolomite contents. The Kellerville also exhibits an abundance of Pennsylvanian lithologies in the very coarse sand through cobble size particles (Hallberg, 1980b). The Alburnett Formation has a much higher limestone to dolomite ratio (almost always greater than 0.40) than the Kellerville Formation.

The differentiation between the Wolf Creek and Alburnett Formation tills is difficult in the field without the assistance of stratigraphic boundaries, but mineralogical characteristics can be used. The most useful characteristic to distinguish between the Wolf Creek and Alburnett formations is the clay mineralogy. The Wolf Creek Formation has higher expandable clay percentages, averaging around 62%; whereas the Alburnett Formation has lower expandable clay percentages, near 43% (Hallberg, 1980a).

**Regional Extent and Thickness:** The Alburnett Formation has a wide range of thickness. In some areas it is completely absent and in others may reach a substantial thickness where its deposits fill in and bury deep bedrock channels. In these areas it has been identified to reach thicknesses of 220-250 feet.

**Origin:** The Alburnett Formation consists of multiple undifferentiated members associated with several Pre-Illinoian ice advances. Based on the physical properties of the Alburnett Formation deposits (massive structure, high density, uniform texture) it is likely a basal, or subglacial, till facies.

**Age:** The Alburnett Formation represents the oldest of the Pre-Illinoian glaciations. Pre-Illinoian deposits in Iowa range from older than 2.2 million years to approximately 500,000 years ago based on volcanic ash dates in western Iowa (Easterbrook and Boellstorff, 1984; Hallberg, 1986). Paleomagnetic studies of the Alburnett Formation in east-central Iowa indicate that these deposits have reversed polarity and are therefore older than 790,000 years (Baker and Stewart, 1984).

## **ACKNOWLEDGEMENTS**

Recognized for contributions to map's production: Andrew B. Asell, Pete Kollasch, Bill Bunker, Dave Pals, Heather Buresh, Katie Foreman, Mary Pat Heitman, and Lois Bair. Drilling was provided under contract with Aquadrill of Iowa City, Iowa; a special thanks to Diane Edberg, and drilling crew members who worked at times in challenging drilling conditions. A special thanks to the following individuals who graciously allowed access to their land for drilling: Dean Arp, Bob Dahms, and Steve Marten. And, a thank you to the Iowa Department of Natural Resources for allowing access on public lands for drilling. Also, thanks to the Scott County Engineer's Office for providing access to all of their boring logs for this mapping project.

## REFERENCES

- Autin, W.J., and Bettis, E.A., III, 1991, Quaternary geology of Mud Creek, eastern Iowa: Report Submitted to the Louisiana State University Board of Regents, LaSER Program, Baton Rouge, 102p.
- Baker, J.L., and Stewart, R.A., 1984, Paleomagnetic study of glacial deposits at Conklin Quarry and other locations in southeast Iowa, *in* Bunker, B.J., and Hallberg, G.R., eds., Underburden-Overburden, an examination of Paleozoic and Quaternary strata at the Conklin Quarry near Iowa City: Geological Society of Iowa Guidebook 41, p. 63-69.
- Baker, R.G., Bettis, E.A. III, Schwert, D.P., Horton, D.G., Chumbley, C.A., Gonzalez, L.A., and Reagan, M.K., Holocene Paleoenvironments of Northeast Iowa, Ecological Monographs, Vol. 66, No. 2 May 1996: Ecological Society of America, p. 203-234.
- Bettis, E.A., III, 1989, Late Quaternary history of the Iowa River Valley in the Coralville Lake area *in* Plocher, O.W., Geologic Reconnaissance of the Coralville Lake area: Geological Society of Iowa Guidebook 51, p. 93-100.
- Bettis, E.A., III, 1990, Holocene alluvial stratigraphy of western Iowa, *in* Bettis, E.A., III, ed., Holocene alluvial stratigraphy and selected aspects of the Quaternary history of western Iowa: Midwest Friends of the Pleistocene Field Trip Guidebook, p. 1-72.
- Bettis, E.A., III, 1995, The Holocene Stratigraphic Record of Entrenched Stream Systems in Thick Loess Regions of the Mississippi River Basin: University of Iowa Department of Geology, Iowa City, unpublished Ph.D. thesis, 149 p.
- Bettis, E.A., III, Baker, R.G., Green, W., Whelan, M.K., and Benn, D.W., 1992, Late Wisconsinan and Holocene alluvial stratigraphy, paleoecology, and archeological geology of east-central Iowa: Iowa Department of Natural Resources-Geological Survey Bureau Guidebook Series No. 12, 82 p.
- Bettis, E.A. III, Hajic, E.R., and Quade, D.J., 1999, Geologic mapping of large valleys in glaciated regions: The use of landform and landscape sediment assemblages for multi-use maps: Geologic Society of America Abstracts with Programs, 33<sup>rd</sup> Annual Meeting North-Central Section April 1999, Champaign, Illinois, vol. 31 no. 5, abstract no. 04164.
- Boellstorff, J., 1978a, North American Pleistocene Stages reconsidered in light of probable Pliocene-Pleistocene continental glaciation: *Science*, v. 202 p.305-307.
- Boellstorff, J., 1978b, Chronology of some late Cenozoic deposits from the central United States and the ice ages: *Transactions of the Nebraska Academy of Science*, v. 6, p. 35-49.
- Daniels, R.B, Rubin, M., and Simonson, G.H., 1963, Alluvial chronology of the Thompson Creek Watershed, Harrison County, Iowa: *American Journal of Science*, v. 261, p. 473-484.
- Easterbrook, D.J., and Boellstorff, J., 1984, Paleomagnetism and chronology of Early Pleistocene tills in the central United States *in* Mahaney, W.C., ed., *Correlation of Quaternary Chronologies*, p. 73-90.
- Esling, S.P., 1984, Quaternary stratigraphy of the lower Iowa and Cedar River valleys, southeast Iowa: University of Iowa, Iowa City, unpublished PhD Dissertation, 451p.

- Eyles, N. and Menzies, J., 1983, The subglacial landsystem, *in* Eyles, N. ed., *Glacial geology-An introduction for engineers and earth scientists*: Oxford, Pergamon, p. 19-70.
- Frye, J.C., Willman, H.B., and Glass, H.D., 1964, Cretaceous deposits and the Illinoian glacial boundary in western Illinois: *Illinois Geological Survey Circular 364*, 28p.
- Frye, J.C., Glass, H.D., Kempton, J.P., and Willman, H.B., 1969, Glacial tills of northwestern Illinois: *Illinois Geological Survey Circular 437*, 47p.
- Hallberg, G.R., 1980a, Pleistocene stratigraphy in east-central Iowa: *Iowa Geological Survey Technical Information Series 10*, 168p.
- Hallberg, G.R., 1980b, Illinoian and Pre-Illinoian stratigraphy of southeast Iowa and adjacent Illinois: *Iowa Geological Survey Technical Information Series 11*, 206p.
- Hallberg, G.R., 1986, Pre-Wisconsin glacial stratigraphy of the central plains region in Iowa, Nebraska, Kansas, and Missouri: *in* Richmond, G.M. and Fullerton, D.S., eds., *Quaternary Glaciations in the United States of America, Report of the International Correlation Programme-Project 24*: *in* Sibrava, V., Bowen, D.Q., and Richmond, G.M., eds., *Quaternary Science Reviews, Quaternary Glaciations in the Northern Hemisphere*, v. 5, p. 11-15.
- Hallberg, G.R., Fenton, T.C., Miller, G.A., and Lutenegeger, A.J., 1978, The Iowan erosion surface: an old story, an important lesson, and some new wrinkles: *in* Anderson, R., ed., *42<sup>nd</sup> Annual Tri-State Geological Field Conference Guidebook*, p. 2-1 to 2-94.
- Hallberg, G.R., Hoyer, B.E., and Miller, G.A., 1974, The geology and paleopedology of the Cherokee Sewer site *in* *The Cherokee Sewer site (13CK405): a preliminary report of a stratified Paleo-Indian/Archaic site in northwestern Iowa*: *Journal of the Iowa Archaeological Society*, v. 21, p. 17-49.
- Hallberg, G.R., and Baker, R.G., 1980, Reevaluation of the Yarmouth type area, *in* Hallberg, G.R., ed., *Illinoian and Pre-Illinoian stratigraphy of southeast Iowa and adjacent Illinois*: *Iowa Geological Survey Technical Information Series 11*, p. 111-150.
- Hallberg, G.R., Wollenhaupt, N.C., and Wickham, J.T., 1980, Pre-Wisconsinan stratigraphy in southeast Iowa, *in* Hallberg, G.R., ed., 1-110.
- Hallberg, G.R., Lineback, J.A., Mickelson, D.M., Knox, J.C., Goebel, J.E., Hobbs, H.C., Whitfield, J.W., Ward, R.A., Boellstorf, J.D., and Swinehart, J.B., 1991, Quaternary geologic map of the Des Moines 4° x 6° quadrangle, United States: U.S. Geological Survey, *Miscellaneous Investigations Series*, Map I-1420, 1:100,000 scale map sheet.
- Hoyer, B.E., 1980a, The geology of the Cherokee Sewer site, *in* Anderson, D.C., and Semken, H.A., Jr. eds. *The Cherokee Excavations: Holocene Ecology and Human Adaptions in Northwestern Iowa*: Academic Press, New York, p. 21-66.
- Hoyer, B.E., 1980b, Geomorphic history of the Little Sioux River Valley: *Geological Society of Iowa Guidebook 34*, 94 p.
- Johnson, W.H., Hansel, A.K., Bettis, E.A., Karrow, P.F., Larson, G.J., Lowell, T.V., and Schneider, A.F., 1997, Late Quaternary temporal and event classifications, Great Lakes region, North America: *Quaternary Research*, v. 47, p. 1-12.

- Kay, G.F., and Graham, J.B., 1943, The Illinoian and post-Illinoian Pleistocene geology of Iowa: Iowa Geological Survey 1940 Annual Report, v. 38, p. 1-262.
- Kemmis, T.J., Bettis, E.A. III, and Hallberg, G.R., 1992, Quaternary geology of Conklin Quarry: Iowa Geological Survey Guidebook Series No. 13, 41 p.
- Leighton, M.M., and Willman, H.B., 1950, Loess formations of the Mississippi Valley: Journal of Geology, v. 58, no. 6, p. 599-623.
- Leverett, F., 1898, The weathered zone (Yarmouth) between the Illinoian and Kansan till sheets: Journal of Geology, v. 6, p. 238-243 and Iowa Academy of Science Proceedings, v. 5, p. 81-86.
- Leverett, F., 1899, The Illinois glacial lobe: U.S. Geological Survey, Monograph 38, 817p.
- Lineback, J.A., 1979, The status of the Illinoian glacial stage: Midwest Friends of the Pleistocene 26<sup>th</sup> Field Conference, Illinois State Geological Survey Guidebook 13, p. 69-78.
- Lineback, J.A., 1980, The Glasford Formation of western Illinois *in* Hallberg, G.R., Illinoian and Pre-Illinoian stratigraphy of southeast Iowa and adjacent Illinois: Iowa Geological Survey Technical Information Series 11, p. 181-184.
- Prior, J.C., 1976, Landforms of Iowa: Iowa City, University of Iowa Press, 154p.
- Quade, D.J., Bettis, E.A. III, and Giglierano, J.D., 2004, Surficial geologic materials of the McCausland Quadrangle, Open File Map 04-5, 1:24,000 scale map sheet.
- Ruhe, R.V., 1969, Quaternary Landscapes in Iowa: Ames, Iowa State University Press, 255p.
- Ruhe, R.V., Dietz, W.P., Fenton, T.E., and Hall, G.F., 1968, Iowan drift problem, northeastern Iowa: Iowa Geological Survey Report of Investigations 7, 10p.
- Sugden, D.E., and John, B.S., 1976, Glaciers and Landscape: New York, John Wiley & Sons, 376 p.
- Wanless, H.R., 1957, Geology and mineral resources of the Beardstown, Glasford, Havana, and Vermont Quadrangles: Illinois Geological Survey Bulletin 82, 233 p.
- Wickham, J.T., 1980, Status of the Kellerville Till Member in western Illinois, *in* Hallberg, G.R., ed., Illinoian and Pre-Illinoian stratigraphy of southeast Iowa and adjacent Illinois: Iowa Geological Survey Technical Information Series 11, p. 151-180.
- Willman, H.B. and Frye, J.C., 1970, Pleistocene stratigraphy of Illinois: Illinois State Geological Survey Bulletin, 94, 204p.
- Willman, H.B., Glass, H.D., and Frye, J.C., 1963, Mineralogy of glacial tills and their weathering profiles in Illinois, Part I- Glacial tills: Illinois Geological Survey Circular 347, 55p.