

Bedrock Geology of the Northwood (Iowa) 7.5' Quadrangle

BEDROCK GEOLOGY OF THE NORTHWOOD 7.5' QUADRANGLE, WORTH COUNTY, IOWA

Iowa Geological and Water Survey
Open File Map OFM-11-02
September 2011

prepared by
Huihao Liu, Robert McKay, Robert Rowden, Deborah Quade, Stephanie Tassier-Sarino,
and James Gagliano
Iowa Geological and Water Survey, Iowa City, Iowa



Iowa Department of Natural Resources, Roger Landis, Director
Iowa Geological and Water Survey, Robert D. Liles, State Geologist

Supported in part by the U.S. Geological Survey
Cooperative Agreement Number G55410621
National Cooperative Geologic Mapping Program (NCGMP)

ACKNOWLEDGMENTS

Special thanks to Keith Brown and Larry Foley for allowing us access to their property, and to Fulkerson LLC, Water Commission Co., T.E. Brown LLC, Trankin & Son Inc., Shell Rock, Inc., and the Worth County Corporation for their assistance in the field. We also thank the Iowa Geological and Water Survey (IGWS) for providing us with the geologic map of the Northwood 7.5' quadrangle. We also thank the IGWS for providing us with the geologic map of the Northwood 7.5' quadrangle. We also thank the IGWS for providing us with the geologic map of the Northwood 7.5' quadrangle.

Introduction to the Bedrock Geology of the Northwood 7.5' Quadrangle

The Northwood 7.5' Quadrangle mostly within Worth County, Iowa, is located near the border area of the Des Moines Lobe landform region, which was the last area covered by Quaternary glacial drift in Iowa, and the Iowan Surface landform region, which was modified by various episodes of erosion before Wisconsinan glacial events (Prior, 1991).

Most of the Northwood quad is covered by Quaternary deposits with a maximum thickness of about 35 m (115 ft), but a few bedrock outcrops were found along the Shell Rock River. Quarries in the southern part of the quad provided significant information for the regional bedrock stratigraphic study. Subsurface information was also derived from the analysis of water well materials collected by Iowa Geological and Water Survey (IGWS) and stored in the GEOSAM database of IGWS. Bedrock information from more than 85 private and public wells within the quad was studied and used for the bedrock geologic mapping.

Middle and lower Upper Devonian rocks form the major bedrock surface and upper bedrock aquifer in the mapping area. This area is within the northern region of the Devonian Iowa Basin. The stratigraphy of this basin has been intensively studied by IGWS staff (e.g., Belanski, 1927, 1928; Koch, 1970) and re-studied and correlated by Witzke and Bunker (1984), Bunker and others (1986), Witzke and others (1988), Anderson and Bunker (1998), Groves and others (2008), etc. The stratigraphic nomenclature and correlation for this map follow the stratigraphic framework proposed by Witzke and others (1988).

The youngest bedrock unit within the quad is the Cretaceous Windrow Formation, which usually occurs as iron-rich reddish erosional outliers a few meters thick in north-central Iowa (Witzke et al., 2010). The Devonian rocks are dominated by carbonates varying between limestone and dolomite, accompanied with minor shale. Based on lithologic features and fossils, the Devonian bedrock in the mapping area can be subdivided into, in descending order, the Shell Rock, Lithograph City, and Coralville formations. The Shell Rock Formation occurs in the southern part of the quad and is characterized by fossiliferous and stromatopore-rich carbonates. Shaly carbonates are also common in the middle portion of the Shell Rock Formation. The underlying Lithograph City Formation is usually represented by laminated lithographic and sublithographic limestone and dolomite and is the dominant bedrock unit in the quad. Locally, a meter-thick fossiliferous and stromatopore-rich facies occurs near the middle of the Lithograph City Formation. The Coralville Formation is characterized by limestone dolomitic limestone, and dolomite, sometimes argillaceous. It forms the bedrock surface in bedrock valleys along northern border of the quad.

Cited References

Anderson, R.R., and Bunker, B.J. (eds), 1998. Fossil shells, glacial swells, and drainage wells: the geology of the Mason City, Iowa, area. *Geol. Soc. of Iowa, Guidebook No. 65*.
Belanski, C.H., 1927. The Shellrock Stage of the Devonian. *American Midland Naturalist*, v. 10, p. 316-370.
Belanski, C.H., 1928. The Shellrock Stage of the Devonian. Description of some typical fossils of the Shellrock Stage. *American Midland Naturalist*, v. 11, p. 165-212.
Bunker, B.J., Witzke, B.J., and Day, J.E., 1986. Upper Cedar Valley Stratigraphy, North-Central Iowa, Lithograph City Formation. *Geol. Soc. of Iowa, Guidebook No. 44*.
Groves, J.R., Walters, J.C., and Day, J. (eds), 2008. Carbonate platform facies and faunas of the Middle and Upper Devonian Cedar Valley Group and Lime Creek Formation, northern Iowa. *IGS Guidebook No. 28*.
Koch, D.L., 1970. Stratigraphy of the Upper Devonian Shell Rock Formation of north-central Iowa. *IGS Report of Investigations 10*, the state of Iowa, 123 p.
Prior, J.C., 1991. Landforms of Iowa. Univ. of Iowa Press, Iowa City, 154 p.
Witzke, B.J., Anderson, R.R., and Pope, J.P., 2010. Bedrock geologic map of Iowa, 1:500,000. IGS Open File Map OFM-2010-01.
Witzke, B.J. and Bunker, B.J., 1984. Devonian stratigraphy of north-central Iowa. *IGS Open File Report 84-2*, p. 107-149.
Witzke, B.J., Bunker, B.J., and Rogers, F.S., 1988. Eifelian through lower Frasnian stratigraphy and deposition in the Iowa area, central midcontinent, U.S.A. in McMillan, N.J., Embry, A.F., and Glass, D.J. (eds), *Devonian of the World*. Canadian Soc. Of Petroleum Geologists, Memoir 14, vol. 1, p. 221-250.

LEGEND

CENOZOIC

QUATERNARY SYSTEM

Qu

Qu - Undifferentiated unconsolidated sediment. Consists of loamy silts developed in loess, glacial till, and colluvium of variable thickness, and alluvial clay, silt, sand, and gravel. Total thickness can be up to 35 m (115 ft) in the quad. This unit is shown only on the cross-section, not on the map.

MESOZOIC

CRETACEOUS SYSTEM

Kw

Kw - Sandstone, Mudsone, and Siderite Pellets (Windrow Formation) "Mid"-Cretaceous. This map unit occurs as erosional outliers and is only found occasionally in well materials in the mapping area. The formation is characterized by reddish shaly sandstone and mudstone or siderite pellets. Its thickness is variable, but usually less than 6 m (20 ft).

PALEOZOIC

DEVONIAN SYSTEM

Dsr

Dsr - Limestone, Dolomite, and Shale (Shell Rock Formation) Upper Devonian. This map unit usually has a thickness of 12 to 18 m (40-60 ft) and occurs in the southern part of the quad. The unit is characterized by fossiliferous carbonates with some grey to light green shale. Layers contain an abundant sub-spherical and tubular stromatopores commonly occur in the lower part of the unit. Brachiopods, bryozoans, corals, and trilobites are abundant in some intervals.

Dlgc

Dlgc - Dolomite, Limestone, and Shale (Lithograph City Formation) Middle to Upper Devonian. This map unit forms the major uppermost bedrock in the quad, with a maximum thickness of up to 3.5 m (110 ft). This unit consists of dolomite and dolomitic limestone, partially characterized by interbeds of laminated lithographic and sublithographic limestone and dolomitic limestone in part argillaceous or with little shale. "Bedrock" or carbonate common. Some intervals are fossiliferous and stromatopore-rich.

Dcv

Dcv - Limestone and Dolomite (Coralville Formation) Middle Devonian. The thickness of this map unit varies between 10 and 18 m (35-60 ft) and is dominated by limestone, dolomitic limestone, and dolomite, in part laminated and argillaceous. Brachiopods and corals usually occur in the limestone facies.

Dlc

Dlc - Dolomite and Limestone (Little Cedar Formation) Middle Devonian. The thickness of this map unit ranges from 27 to 35 m (90-115 ft) in the study area. The unit is dominated by slightly argillaceous to argillaceous dolomitic and dolomitic limestone, usually wavy and partially laminated and/or cherty. This unit is commonly fossiliferous, and brachiopods are especially abundant in the lower portion. This unit is shown only in the cross-section, not on the map.

•



Drill Holes

x


Outcrops

Correlation of Map Units

AGE (Ma)	SYSTEM	SERIES	STAGE	MAP UNIT
2.58	QUATERNARY			Qu
145.5	CRETACEOUS			Kw
385	DEVONIAN	Upper	Frasnian	Dsr
390			Givetian	Dlgc
395			Eifelian	Dcv
		Middle		Dlc



GEOLOGIC CROSS-SECTION A-B



Base map from USGS Northwood 7.5' Digital Raster Graphic (DRG) file 010608101 which was scanned from the Northwood 7.5' Topographic Quadrangle map published by US Geological Survey in 1973. Topographic contour lines are based on the 1973 map. The map and cross-section are based on interpretations of the best available information at the time of mapping. Map interpretations are not a substitute for detailed site-specific studies.